

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/02, 5/10</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 99/14316</b> <b>(43) International Publication Date:</b> 25 March 1999 (25.03.99)
<b>(21) International Application Number:</b> PCT/US98/19571 <b>(22) International Filing Date:</b> 18 September 1998 (18.09.98)  <b>(30) Priority Data:</b> 60/059,234 18 September 1997 (18.09.97) US  <b>(71) Applicant (for all designated States except US):</b> AGRITOPE, INC. [US/US]; 16160 S.W. Upper Boones Ferry Road, Portland, OR 97224-7744 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> BESTWICK, Richard, K. [US/US]; 6680 S.W. Canby, Portland, OR 97223 (US). KELLOGG, Jill, A. [US/US]; 6680 S.W. Canby, Portland, OR 97223 (US).  <b>(74) Agents:</b> JUDGE, Linda, R. et al.; Dehlinger & Associates, P.O. Box 60850, Palo Alto, CA 94306-0850 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> SYNTHETIC HYBRID PLANT PROMOTER  <b>(57) Abstract</b>  The present invention is directed to a synthetic hybrid promoter composed of polynucleotide segments derived from the E8 and E4 gene promoters. The hybrid promoter is capable of providing high-level expression of heterologous genes, particularly in transformed fruit. DNA constructs containing the E8-E4 hybrid promoter operably linked to an exemplary heterologous SAMase gene are effective in conferring a delayed ripening phenotype to transformed fruit.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## SYNTHETIC HYBRID PLANT PROMOTER

### Field of the Invention

The present invention relates to a synthetic E8-E4 hybrid promoter composed of  
5 polynucleotide segments derived from the tomato E8 and tomato E4 genes, and to DNA constructs,  
chimeric genes, vectors, kits, and transformation methods employing the promoter.

### References

- Adams, D.O., and Yang, S.F., *Plant Physiology* 70:117-123 (1977).
- 10 Ausubel, F.M., *et al.*, in CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley and  
Sons, Inc., Media, PA (1992).
- Ayub, R., *et al.*, *Nature Biotechnology* 14:862-866 (1996).
- Balague, C., *et al.*, *Eur. J. Biochem.* 212:27-34 (1993).
- Balazs, E., *et al.*, *Gene* 19(3):239-249 (1982).
- 15 Becker, D., *et al.*, *Plant Mol. Biol.* 20:1195-1197 (1992).
- Bellini, C., *et al.*, *Bio/Technology* 7(5):503-508 (1989).
- Brunke, K.J. and Wilson, S.L., European Patent Publication No. 0 559 603 A2, published  
September 08, 1993.
- Comai, L., and Coning, A.J., U.S. Patent No. 5,187,267, issued 16 February 1993.
- 20 Cordes, S., *et al.*, *The Plant Cell* 1:1025-1034 (1989).
- Coupe, S.A. and Deikman, J., *The Plant Journal* 11(6):1207-1218 (1997).
- Dayhoff, M.O., in ATLAS OF PROTEIN SEQUENCE AND STRUCTURE Vol. 5, National  
Biomedical Research Foundation, pp. 101-110, and Supp. 2, Vol. 5, pp. 1-10 (1972).
- Deikman, J., *et al.*, *Plant Physiol.* 100(4):2013-2017 (1992).
- 25 Deikman, J. and Fischer, R.L., *EMBO J.* 7:3315-3320 (1988).
- Dong, J.Z., *et al.*, *Bio/Technology* 9:858-863 (1991).
- Fang, G., and Grumet, R., *Plant Cell Rep.* 9:160-164 (1990).
- Fraley, R., *et al.*, U.S. Patent No. 5,352,605, issued on October 4, 1994.
- Gonsalves, C., *et al.*, *J. Amer. Soc. Hort. Sci.* 119:345-355 (1994).
- 30 Good, X., *et al.*, *Plant Mol. Biol.* 26:781-790 (1994).
- Guilley, H., *et al.*, *Cell* 30(3):763-773 (1982).
- Hamilton, A.J., *et al.*, *Nature* 346:284-287 (1990).
- Hood, E., *et al.*, *Transgenic Research* 2:208-218 (1993).

- Hooykaas, P.J.J., and Schilperoot, R.A., in TRENDS IN BIOCHEMICAL SCIENCES, International Union of Biochemistry and Elsevier Science Publishers, v. 10(8):307-309 (1985).
- Houck, C.M. and Pear, J.R., U.S. Patent No. 4,943,674, issued 24 July 1990.
- Jefferson, R.A., *Plant Mol. Biol. Rep.* 5:387 (1987b).
- 5 Jefferson, R.A., *et al.*, *EMBO J.* 6:3901 (1987a).
- Jefferson, R.A., *et al.*, *EMBO J.* 6:3901 (1987a).
- Jefferson, R.A., *Plant Mol. Biol. Rep.* 5:387 (1987b).
- Klee, H.J., *et al.*, *Plant Cell* 3:1187-1193 (1991).
- Klein, T.M., *et al.*, *PNAS USA* 85(22):8502-8505 (1988).
- 10 Knessl, M.L. and Deikman, J., *Plant Physiology* 112:537-547 (1996).
- Kramer, M.G., *et al.*, in BIOLOGY & BIOTECHNOLOGY OF THE PLANT HORMONE ETHYLENE Kluwer Academic Publishers, The Netherlands (1996).
- Laemmli, E.K., *Nature* 227:680-685 (1970).
- Leisner, S.M., and Gelvin, S.B., *Proc. Natl. Acad. Sci. USA* 85(8):2553-2557 (1988).
- 15 Lin, E., *et al.*, *Plant Mol. Biol.* 23:489-499 (1993).
- Lincoln, J.E., *et al.* *Proc. Natl. Acad. Sci. U.S.A.*, 84:2793-2797 (1987).
- Lincoln, J.E. and Fischer, R.L., *Mol. Gen. Genet.* 212:71-75 (1988).
- Maniatis, T., *et al.*, in MOLECULAR CLONING: A LABORATORY MANUAL, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1982).
- 20 Mariani, C., *et al.*, *Nature* 357:384-387 (1992).
- Mathews, H.V., *et al.*, *In Vitro Cell Dev. Biol.* 28P:172-178 (1992).
- Melchers, L.S., *et al.*, *Plant J.* 5:469-480 (1994).
- Miki, B.L.A., *et al.*, PLANT DNA INFECTIOUS AGENTS (Hohn, T., *et al.*, Eds.) Springer-Verlag, Vienna, Austria, pp. 249-265 (1987).
- 25 Montgomery, J., *et al.*, *Plant Cell* 5:1049-1062 (1993).
- Montgomery, J., *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 90:5939-5943 (1993).
- Mullis, K.B., *et al.*, U.S. Patent No. 4,683,195, issued 28 July 1987.
- Mullis, K.B., U.S. Patent No. 4,683,202, issued 28 July 1987.
- Ni, M., *et al.*, *Plant J.* 7:661-676 (1995).
- 30 Odell, J.T., *et al.*, *Plant Mol Biol* 10(3):263-272 (1988).
- Odell, J.T., *et al.*, *Nature* 313:810-812 (1985).
- Odell, J.T., *et al.*, *J. Cell Biochem. (Suppl. 11B)*:60 (1987).
- Pearson, W.R., *Methods in Enzymol.* 183:63-98 (1990).
- Pearson, W.R., and Lipman, D.J., *PNAS* 85:2444-2448 (1988).

- Penarrubia, L., *et al.*, *Plant Cell* 4:681-687 (1992).
- Picton, S., *et al.*, *Plant Physiology* 103(4):1471-1472 (1993).
- Ponstein, A.S., *et al.*, *Plant Physiol.* 104:109-118 (1994).
- Rogers, S., U.S. Patent No. 5,378,619, issued on January 3, 1995.
- 5 Rogers, S., U.S. Patent 5,034,322, issued on July 23, 1991.
- Saiki, R.K., *et al.*, *Science* 239:487-491 (1988).
- Sambrook, J., *et al.*, in MOLECULAR CLONING: A LABORATORY MANUAL, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, Vol. 2 (1989).
- Sato, T., and Theologis, A., *Proc. Natl. Acad. Sci. USA* 86:6621-6625 (1989).
- 10 Schuch, W., *Euphytica*. 79(3):287-291 (1994).
- Sheehy, R.E., *et al.*, *J. Bact.* 173:5260-5265 (1991).
- Studier, F.W., *et al.*, *J. Virol* 19:136 (1976).
- Tommerup, H., *et al.*, *Eur. Congr. Biotechnol.* 5:916-918 (1990).
- Toubart, P., *et al.*, *Plant J.* 3:367-373 (1992).
- 15 Valles, M.P. and Lasa, J.M., *Plant Cell Rep.* 13:145-148 (1994).
- Van Haaren, M.J.J., *et al.*, *Plant Mol. Bio.* 21:625-640 (1993).
- Woloshuk, C.P., *et al.*, *J. Plant Cell* 3:619-628 (1991).
- Xu, R., *et al.*, *Plant Mol. Biol.* 31:1117-1127 (1996).
- Yoshioka, K., *et al.*, *Jpn. J. Breeding* 42(2):278-285 (1992).
- 20 Zhu, Q., *et al.*, *Plant Cell* 7:1681-1689 (1995).

### **Background of the Invention**

Promoters that regulate gene expression in plants are essential elements of plant genetic engineering. Several examples of promoters useful for the expression of selected genes in plants are  
 25 now available (Zhu, *et al.*, 1995; Ni, *et al.*, 1995).

To be expressed in a cell, a gene must be operably linked to a promoter which is recognized by certain enzymes in the cell. The 5' non-coding regions of a gene (*i.e.*, regions immediately 5' to the coding region), referred to as promoters or transcriptional regulatory regions, initiate transcription of the gene to produce a mRNA transcript. The mRNA is then translated at the ribosomes of the cell  
 30 to yield an encoded polypeptide.

Promoters typically contain from about 500-1500 bases, and can provide regulated expression of genes under their control. Expression of heterologous genes or selected sequences of genes in transgenic plants has typically involved the use of constitutive promoters, *i.e.*, promoters which drive the expression of a product throughout the plant at all times and in most tissues.

Promoters derived from viral genes are useful for expressing selected genes in plants. Examples of such viral genes which have been identified include those found in the caulimovirus family of viruses (a group of double-stranded DNA viruses), and include the Cauliflower Mosaic Virus (CaMV) 35S (Balazs, *et al.*, 1982; Guilley, *et al.*, 1982; Odell, *et al.*, 1985; Odell, *et al.*, 1987; 5 Odell, *et al.*, 1988; Tommerup, *et al.*, 1990; Jefferson, *et al.*, 1987a; Jefferson, 1987b) and CAMV 19S promoters (Fraley, *et al.*, 1994), and the Figwort Mosaic Virus (FMV) (Rogers, 1995) promoter.

Promoters useful for regulating gene expression in plants and obtained from bacterial sources, *e.g.*, *Agrobacterium*-derived promoters, have been identified and isolated. Such promoters include those derived from *Agrobacterium* T-DNA opine synthase genes, and include the nopaline synthase 10 (nos) promoter (Rogers, 1991), the octopine synthase (ocs) promoter (Leisner and Gelvin, 1988) and mannopine synthase (mas) promoter.

Plant promoters (promoters derived from plant sources) effective to provide constitutive expression, are less well known, and include *hsp80*, Heat Shock Protein 80 from cauliflower, (Brunke and Wilson, 1993), and the tomato ubiquitin promoter (Picton, *et al.*, 1993). These promoters can 15 be used to direct the constitutive expression of heterologous nucleic acid sequences in transformed plant tissues.

The methods and results described herein demonstrate the ability to provide tissue and/or stage specific regulation of gene expression in transgenic plants. The tissue and/or stage-specific promoters of the present invention include a region of DNA that regulates transcription of the immediately 20 adjacent (downstream) gene to a specific plant tissue. According to methods of the present invention, heterologous genes are linked to the promoters of the present invention. Exemplary heterologous gene for the transformation of plants include genes whose products are effective to reduce ethylene biosynthesis in specific tissues of those plants, *e.g.* the fruits. An exemplary gene which impacts fruit ripening is the bacteriophage T3 gene encoding S-adenosylmethionine hydrolase (SAMase), Kramer, 25 *et al.*, (1996); Good, *et al.*, (1994).

A limited number of inducible and/or tissue specific promoters are known. Promoters that provide fruit-specific expression include the E4 and E8 promoter from tomato (Cordes, *et al.*, 1989; Bestwick, *et al.*, 1995), and corresponding promoters from other plants which have substantially the same biological activity. The tomato E4 promoter is both stage and tissue specific (Cordes, *et al.*, 30 1989), and typically, E4 mRNA is abundant in ripening tomato fruit and is not detected in leaf, root, stem, or unripe fruit. The E4 and E8 promoters are also ethylene responsive promoters, *i.e.*, transcription of genes placed under their control is activated by ethylene. The E4 promoter requires ethylene for activation, while the E8 promoter is controlled by ethylene as well as by separate developmental signals (Deikman *et al.*, 1997). Another fruit-specific promoter is the tomato 2AII 35 gene promoter. It has been demonstrated that nucleic acid sequences placed under the regulatory

control of the 5' non-coding region of the tomato 2All gene (Van Haaren) are preferentially transcribed in developing fruit tissue. Fruit specific regulation of the kiwifruit actinidin promoter has been reported to be conserved in transgenic petunia plants (Lin, *et al.*, 1993).

At present, a relatively small number of plant promoters, particularly constitutive plant promoters, have been identified. The use of such promoters in plant genetic engineering has been rather limited to date, since gene expression in plants is, for the most part, typically tissue, developmentally, or environmentally-regulated.

A need exists for tissue and developmental stage specific promoters that are functional in plant cells, and which are capable of providing high level expression of heterologous genes.

10

### Summary of the Invention

The present invention is directed to a synthetic promoter composed of a combination of cis-acting elements derived from the transcriptional regulatory sequences of the E8 and E4 genes, as exemplified by tomato. The synthetic hybrid promoter allows high-level, fruit specific expression of nucleic acid sequences placed under its control.

In one aspect, the invention provides a DNA construct which contains a DNA coding sequence under the transcriptional control of a hybrid E8-E4 promoter. The DNA coding sequence is typically heterologous to the hybrid promoter and is operably linked to the promoter to enable expression of the product. Exemplary products include, but are not limited to S-adenosylmethionine hydrolase (SAMase), amino-cyclopropane-1-carboxylic acid (ACC) deaminase, ACC oxidase antisense molecule, ACC synthase antisense molecule, ACC oxidase cosuppression molecule, ACC synthase cosuppression molecule, thaumatin, sucrose phosphate synthase and lycopene cyclase.

The E8-E4 hybrid promoter of the invention is composed of a polynucleotide segment derived from an E8 gene promoter which is fused to a polynucleotide segment derived from an E4 gene promoter positioned downstream of the E8 promoter segment. The E8 promoter-derived polynucleotide segment includes at least 30 contiguous nucleotides selected from the region extending from nucleotide positions -2257 to -847 of the tomato E8 promoter which corresponds to approximately nucleotides 1 to 1411 of SEQ ID NO:7, or the functional equivalent thereof. The polynucleotide segment derived from an E4 gene promoter includes at least 200 contiguous nucleotides selected from the region extending from nucleotide positions -1150 to +16 of the tomato E4 promoter which corresponds to approximately nucleotides 271 to 1437 of SEQ ID NO:8, or the functional equivalent thereof. The particular combination of E8 and E4 polynucleotide segments produces a hybrid promoter which is effective to drive expression of a heterologous gene (*e.g.*, a reporter gene) at a level of at least about 75-300% of the expression level obtained using either an unmodified E4 or E8 gene promoter.

In one embodiment of the hybrid promoter, the nucleotide sequence of the E8 promoter segment corresponds to nucleotides -1529 to -847 of the tomato E8 promoter, or the functional equivalent thereof, which corresponds to approximately nucleotides 3 to 686 of SEQ ID NO:6 and nucleotides 729 to 1411 of SEQ ID NO:7. The hybrid promoter also contains an E4 promoter  
5 segment corresponding to nucleotides -315 to +16 of the tomato E4 promoter, or the functional equivalent thereof, which corresponds to approximately nucleotides 693 to 1023 of SEQ ID 6 or nucleotides 1107 to 1437 of SEQ ID NO:8, referred to herein as the "short E8-E4 hybrid promoter".

In another embodiment of the hybrid promoter, the nucleotide sequence of the E8 promoter segment corresponds to nucleotides -2257 to -1103 of the tomato E8 promoter, or the functional  
10 equivalent thereof, which corresponds to approximately nucleotides 1 to 1160 of SEQ ID NO:1 and nucleotides 1 to 1156 of SEQ ID NO:7. The hybrid promoter also contains an E4 promoter segment corresponding to nucleotides -1150 to +16 of the tomato E4 promoter, or the functional equivalent thereof, which corresponds to approximately nucleotides 1157 to 2323 of SEQ ID 1 or nucleotides 271 to 1437 of SEQ ID NO:8, designated herein as the "long E8-E4 promoter".

15 In one respect, the E8-E4 hybrid promoter of the present invention can be used to reduce ethylene production in transformed fruit cells, to thereby alter the ripening phenotype of transgenic fruit composed of such fruit cells.

In another embodiment, the DNA sequence can correspond to a pathogenesis related gene, such as polygalacturonase inhibiting protein (PGIP), glucanase and chitinase.

20 The present invention also includes the use of any of the above chimeric gene constructs to generate a plant transformation vector. Such vectors can be used in any plant cell transformation method, including *Agrobacterium*-based methods, electroporation, microinjection, and microprojectile bombardment. These vectors may form part of a plant transformation kit. Other components of the kit may include, but are not limited to, reagents useful for plant cell transformation.

25 In another embodiment, the invention includes a plant cell, plant tissue, transgenic plant, fruit cell, whole fruit, seeds or calli containing any of the above-described chimeric genes and the corresponding expressed gene products. Preferred plants are dicots such as melon, strawberry, raspberry, and tomato, and particularly *Cucumis sp* (e.g., *Cucumis melo*).

In another aspect of the present invention, the hybrid promoters described herein are employed  
30 in a method for delaying ripening of fruit from a fruit-bearing plant. In this method, a transgenic plant containing the chimeric gene of the present invention is grown to produce a transgenic plant bearing fruit. In one particular embodiment, the chimeric gene encodes a product capable of reducing ethylene biosynthesis when expressed in plant cells (e.g., S-adenosyl-methionine hydrolase, aminocyclopropane-1-carboxylic acid (ACC) deaminase, ACC oxidase antisense molecule, ACC synthase  
35 antisense molecule, ACC oxidase cosuppression molecule, ACC synthase cosuppression molecule).

Fruit produced by these transgenic plants have a modified ripening phenotype. A modified ripening phenotype refers to an alteration of the rate of ripening (*e.g.*, an increased time course, or delay of ripening) of a transgenic fruit relative to corresponding (*i.e.*, non-transgenic) wild-type fruit.

Further, the invention includes a method for producing a transgenic plant such as a fruit-bearing plant. In this method, the chimeric gene of the present invention, typically carried in an expression vector allowing selection in plant cells, is introduced into progenitor cells of selected plant. These progenitor cells are then grown to produce a transgenic plant bearing fruit.

Yet another aspect of the invention is directed to a method for conferring enhanced expression activity to an E4 promoter. In the method, a polynucleotide segment of at least 30 contiguous nucleotides selected from the region extending from nucleotide positions -2257 to -847 of the tomato E8 promoter which corresponds to approximately nucleotides 1 to 1411 of SEQ ID NO:7, or the functional equivalent thereof, is fused in an upstream orientation to an E4 promoter polynucleotide segment of at least 200 contiguous nucleotides selected from the region extending from nucleotide positions -1150 to +16 of the tomato E4 promoter which corresponds to approximately nucleotides 271 to 1437 of SEQ ID NO:8, or a functional equivalent thereof, to form a hybrid E8-E4 promoter capable of regulating expression of a heterologous gene operably linked thereto. The hybrid promoter is effective to drive expression of the heterologous gene to a greater degree than the unmodified E4 promoter, and preferably at a level of at least about 75-300% of the expression level obtained by using an unmodified E4 gene promoter. The hybrid promoter is also ethylene inducible and is capable of directing fruit-specific expression.

These and other objects and features of the invention will become more fully apparent when the following detailed description is read in conjunction with the accompanying figures and examples.

## **Brief Description of the Figures**

Fig. 1 presents a schematic description of the details of the construction of vector pAG-1762 (14.10 kb);

Figs. 2A and 2B correspond to the nucleotide sequence of the long (large) E8-E4 promoter of the present invention (SEQ ID NO:1);

Fig. 3 presents a schematic representation of the details of the construction of an intermediate vector pAG-7162 (14.80 kb);

Fig. 4 presents a schematic representation of the details of the construction of an intermediate transfer vector pAG-126 (4186 bp);

Fig. 5 presents a schematic representation of plasmid pAG-7182 (13.50 kb);

Fig. 6 is a computer-generated image of a Western blot indicating SAMase expression in ripe melons transformed with binary vectors pAG7142 (tomato E4::SAMase) and pAG7152 (tomato E8::SAMase);

Fig. 7 is a computer-generated image of a Western blot indicating SAMase expression in ripe melons transformed with binary vectors pAG7162 (long or large tomato E8-E4::SAMase) and pAG7182 (short or small tomato E8-E4::SAMase);

Fig. 8 is a bar graph representing ethylene biosynthesis profiles over a 4-day period for three different transgenic events (pAG-7142 and control; pAG-7152 and control; and pAG-7162 and control);

Fig. 9 is a schematic representation of the construction of vector pAG-7152 containing the tomato E4 promoter (13.7 kb);

Fig. 10 presents the nucleotide sequence of the short (small) E8-E4 promoter of the present invention (SEQ ID NO:6);

Fig. 11 is a schematic representation of active polynucleotide segments derived from the tomato E8 and E4 promoters which were utilized to prepare the hybrid promoters of the present invention, and

Fig. 12 is a schematic representation of the construction of vector pAG-7142 containing the tomato E8 promoter (14.5 kb).

## **Detailed Description of the Invention**

### **I. Definitions**

The term "polynucleotide" as used herein refers to a polymeric molecule having a backbone that supports bases capable of hydrogen bonding to typical polynucleotides, where the polymer backbone presents the bases in a manner to permit such hydrogen bonding in a sequence specific fashion between the polymeric molecule and a typical polynucleotide (*e.g.*, single-stranded DNA). Such bases are typically inosine, adenosine, guanosine, cytosine, uracil and thymidine. Polymeric molecules include double and single stranded ribonucleic acids (RNA) and deoxyribonucleic acids (DNA), and may include polymers having backbone modifications such as methylphosphonate linkages.

The term, "recombinant nucleic acid" as used herein refers to nucleic acid, originally formed *in vitro*, in general, by the manipulation of nucleic acid by endonucleases, in a form not normally found in nature.

A "heterologous" DNA coding sequence is a structural coding sequence that is not native to the plant being transformed, or a coding sequence that has been engineered for improved characteristics of its protein product. Heterologous, with respect to the promoter, refers to a coding

sequence that does not exist in nature in the same gene with the promoter to which it is currently attached.

Nucleic acid subunits are referred to herein by their standard base designations; T, thymine; A, adenosine; C, cytosine; G, guanine, U, uracil; variable positions are referred to by standard IUPAC abbreviations: W, A or T/U; R, A or G; S, C or G; K: G or T/U (37 CFR. §1.822).

The nucleic acid may be double stranded, single stranded, or contain portions of both double stranded or single stranded sequence. The depiction of a single strand also defines the sequence of the other strand and thus also includes the complement of the sequence.

A "heterologous" DNA or gene sequence encodes a gene product not normally contiguous or associated with the promoter (*e.g.*, an E8-E4 hybrid promoter adjacent DNA sequences encoding S-adenosylmethionine cleaving enzyme). In the context of the present invention, a heterologous gene is any DNA sequence other than the E8 or E4 gene sequence.

"Regulatable promoter" is any promoter whose activity is affected by a *cis* or *trans* acting factor (*e.g.*, an ethylene-inducible promoter such as the tomato E8 promoter).

"Constitutive promoter" is any promoter that directs RNA production in many or all tissues of a plant transformant at most times.

A "tissue-specific promoter" is any promoter which directs RNA synthesis at higher levels in particular types of cells and tissues (*e.g.*, a fruit specific promoter);

By "promoter" or "promoter segment" (*e.g.*, a tomato E8 or E4 promoter segment) is meant a sequence of DNA that functions in a hybrid promoter disclosed herein to direct transcription of a downstream heterologous gene, and includes promoter or promoter segments derived by means of ligation with operator regions, random or controlled mutagenesis, addition or duplication of enhancer sequences, addition or modification with synthetic linkers, and the like, having promoter activity the functional equivalent of, the E8-E4 hybrid promoter described herein or pertinent regions thereof.

By "plant promoter" is meant a promoter or promoter region (as defined above), which in its native form, is derived from plant genomic DNA. The hybrid promoter of the present invention is a plant promoter.

By an E8 or an E4 gene promoter is meant a promoter obtained from an E8 or E4 gene considered to share sequence identity with the tomato E8 or E4 gene sequences described herein, or a particular region or regions thereof, or from a gene having at least about 70%, preferably about 80%, more preferably about 85%, even more preferably about 90% sequence identity over a length of polynucleotide sequence corresponding to the tomato E8 or tomato E4 gene sequences described herein.

Alternatively, an E4 or E8 gene promoter is obtained from the gene encoding an E8 or E4 protein wherein the amino acid sequence of the E4 or E8 protein has at least about 70%, preferably

about 80%, more preferably about 85%, even more preferably about 90% sequence identity over a length of polypeptide sequence corresponding to the tomato E8 or tomato E4 polypeptide sequences of SEQ ID NO:10 and SEQ ID NO:9, respectively.

The term "homology" or "homologue" as used herein refers to the level of identity between two sequences, *i.e.* 70% homology means the same thing as 70% sequence identity as determined by the algorithms described below, and accordingly a homologue of a given sequence has at least about 70%, preferably about 80%, more preferably about 85%, even more preferably about 90% sequence identity over a length of the given sequence.

"Promoter strength" refers to the level of promoter-regulated expression of a heterologous gene in a plant tissue or tissues, relative to a suitable standard (*e.g.*, a hybrid E8-E4 promoter from a particular plant, *e.g.*, tomato, versus either the tomato E8 gene or tomato E4 gene promoter alone). Expression levels can be measured by linking the promoter to a suitable reporter gene such as GUS ( $\beta$ -glucuronidase), dihydrofolate reductase, or *nptII* (neomycin phosphotransferase II). Expression of the reporter gene can be easily measured by fluorometric, spectrophotometric or histochemical assays (Jefferson, *et al.*, 1987a; Jefferson, 1987b).

For the purposes of the present invention, a high level E4/E8 hybrid promoter is one that drives expression of a particular gene, such as a reporter gene, at about 75-300% of the levels obtained with either the non-hybrid E4 or E8 gene promoter derived from the same source.

"Nucleic acid sequence identity" is determined essentially as follows. Two polynucleotide sequences of the same length are considered to be identical to one another, if, when they are aligned using the ALIGN program, over 60%, preferably about 70%, preferably about 80%, more preferably about 85%, even more preferably about 90% sequence are determined to be identical when aligned using the default parameters and the default PAM matrix (Dayhoff, 1972). The ALIGN program is found in the FASTA version 1.7 suite of sequence comparison programs (Pearson and Lipman, 1988; Pearson, 1990; program available from William R. Pearson, Department of Biological Chemistry, Box 440, Jordan Hall, Charlottesville, VA).

"Amino acid sequence identity" with respect to the amino acid sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the native sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Such amino acid sequence identity may be determined using the ALIGN program, with default parameters as described above.

By "functional equivalent" is meant a nucleic acid sequence which corresponds to an active E4 or E8 promoter wherein the E4 or E8 gene product expressed by the active E4 or E8 promoter has at least about 80% amino acid sequence identity with the E4 or E8 gene product having the amino

acid sequence shown in SEQ ID NOs:10 or 9, respectively. The functional equivalent of an E4 or E8 promoter includes, for instance, an E4 or E8 promoter which has a nucleic acid sequence that is the same as part of, but not all of, the nucleic acid sequence of the E4 or E8 promoter and which either retains essentially one or more of the same biological functions or activities as a full length E4 or E8 promoter, *e.g.*, the tomato E4 or E8 promoter, as described herein.

Two nucleic acid fragments are considered to be "selectively hybridizable" to a reference polynucleotide if they are capable of specifically hybridizing to the polynucleotide or variants thereof or of specifically priming a polymerase chain amplification reaction: (i) under moderately stringent hybridization and wash conditions, as described, for example, in Maniatis, *et al.* (1982), pages 320-328, and 382-389; (ii) using reduced stringency wash conditions that allow at most about 25-30% base pair mismatches, for example:  $2 \times$  SSC (contains sodium 3.0 M NaCl and 0.3 M sodium citrate, at pH 7.0), 0.1% sodium dodecyl sulfate (SDS) solution, room temperature twice, 30 minutes each; then  $2 \times$  SSC, 0.1% SDS, 37°C, once, for 30 minutes; then  $2 \times$  SSC, at room temperature twice, for 10 minutes each, or (iii) selecting primers for use in typical polymerase chain reactions (PCR) under standard conditions (for example, in Saiki, *et al.*, 1988), which result in specific amplification of sequences of the desired target sequence or its variants.

As used herein, a "plant cell" refers to any cell derived from a plant, including undifferentiated tissue (*e.g.*, callus) as well as plant seeds, pollen, progagules and embryos.

## 20 II. Inducible Promoters for Expression of Genes in Plants and the Role of Ethylene

The present invention is directed to the applicants' discovery of a hybrid promoter prepared by a combination of regions derived from an E8 and an E4 promoter that is capable of directing high level expression of heterologous genes.

In preparing transgenic plants, a desirable feature is the ability to express an introduced transgene in response to a particular stimulus, or to localize its expression to certain tissues. In an effort to develop new tissue-specific and/or developmentally activated promoters capable of directing high level expression of a heterologous gene in plants, it was discovered that a combination of regions derived from the tomato E8 and E4 promoters, when ligated in a 5' to 3' fashion, were capable of providing such a promoter.

To summarize, the present invention is based upon the surprising discovery of a high-level hybrid E8-E4 promoter which (i) is capable of driving expression of a heterologous gene at significantly greater levels than either the unmodified E8 or E4 promoters alone, (ii) retains the fruit and ripening-specific function of the parent regulatory regions and (iii) is effective in the plant from which the E4 and E8 promoter sequences were derived (*e.g.*, tomato), as well as in other plants (*e.g.*, muskmelon, apple and pear).

The parent promoters from which the hybrid promoter is derived were selected due to a number of features, and in particular, their ability to regulate expression of the SAMase gene in ripening tomato fruit (Lincoln and Fischer, 1988; Lincoln, *et al.*, 1987).

The SAMase gene encodes the enzyme S-adenosyl-methionine hydrolase. The isolation, cloning and sequence of the SAMase gene is described in U.S. Patent No. 5,589,623, and in International PCT Publication No. WO 95/35387. This enzyme, encoded by the *E. coli* bacteriophage T3, hydrolyzes AdoMet to homoserine and MTA. The enzyme is referred to as AdoMet hydrolase (AdoMetase), or by its other name, S-adenosylmethionine cleaving enzyme (SAMase) (Studier, *et al.*). When expressed in plant cells, AdoMetase is effective to "short circuit" a branch of the biosynthetic pathway that produces ethylene, thereby reducing ethylene production in transgenic plants expressing the gene.

The effects of ethylene on plants, whether produced by the plant itself or applied exogenously, are numerous and of considerable commercial importance. Among the diverse physiological effects are leaf abscission, fading and wilting of flowers, leaf yellowing, leaf epinasty, and stimulation of ripening in fruits and vegetables. Of even greater significance is the fact that ethylene promotes senescence (a natural, genetically controlled degenerative process which typically leads to death) in plants, both in selected groups of cells and in whole organs, such as, fruits, leaves, or flowers.

Normally, ethylene production from plant tissue is low. Large quantities of ethylene, however, are produced during ripening and senescence processes, and are also produced following trauma. In fruits and vegetables, the stimulation of ethylene production by cuts or bruises can bear considerably on the effectiveness of storage. Ethylene-induced leaf browning is a common basis for loss in many plants, including lettuce and tobacco. In some tissues, exposure to only a small amount of ethylene may cause an avalanche of ethylene production in adjacent plants or plant tissues such as fresh produce. This autocatalytic effect can be very pronounced and lead to loss of fruit quality during transportation and storage.

Thus, in one aspect, the present invention provides a method to regulate plant cell expression of any gene in a tissue or development stage-specific manner, in particular, genes whose products reduce ethylene synthesis in plant cells, using a hybrid promoter of the type described herein.

Returning now to the tomato E8 and E4 genes from which segments of the exemplary hybrid promoter are derived, the intact parent promoters, the tomato E8 and the tomato E4 promoter, are ethylene inducible (Deikman, *et al.*, 1992; Xu, *et al.*, 1996).

The tomato E4 and E8 promoters may be used to isolate functionally equivalent promoters from other plants. For example, the raspberry E4 promoter may be obtained from a raspberry homologue of the tomato E4 gene. Accordingly, the tomato E4 and E8 promoters can be used to

isolate functionally equivalent promoters or partial sequences thereof from additional other types of plants, and those promoter sequences used to make hybrid E4/E8 promoters.

### III. Construction of a Hybrid E8-E4 Promoter

5       The E8-E4 hybrid promoter contains a combination of nucleotide segments as exemplified by those derived from the tomato E8 and E4 genes. These segments, when combined in a 5'-to-3' fashion, are capable of providing a promoter having certain features, as will be described below.

The component polynucleotide segments of the hybrid promoter were determined on the basis of experiments conducted in support of the invention, as described in Examples 1-7.

10       The E8-E4 hybrid promoter is composed of a polynucleotide segment derived from an E8 gene promoter which is fused to a polynucleotide segment derived from an E4 gene promoter positioned downstream of the E8 promoter segment. The E8 promoter-derived polynucleotide segment preferably includes at least 30 contiguous nucleotides selected from the region extending from nucleotide positions -2257 to -847 of the tomato E8 promoter which corresponds to approximately  
15 nucleotides 1 to 1411 of SEQ ID NO:7, or the functional equivalent thereof. The E8 promoter sequence of SEQ ID NO:7 consists of the E8 promoter described by Deikman and Fischer, 1988 and Deikman, *et al.*, 1992, extended on the 5' end, as described in Example 1. The polynucleotide segment derived from a E4 gene promoter preferably includes at least 200 contiguous nucleotides selected from the region extending from nucleotide positions -1150 to +16 of the tomato E4 promoter  
20 which corresponds to approximately nucleotides 271 to 1437 of SEQ ID NO:8, or the functional equivalent thereof.

The construction of exemplary vectors containing the hybrid promoter is typically carried out as described in Examples 1-6. The sequence of the tomato E8 promoter for use in the present invention is provided in SEQ ID NO:7, and the DNA sequence of regions relevant to the examples  
25 of the present invention are provided in Figs. 2A and 2B (SEQ ID NO:1) and Fig. 10 (SEQ ID NO:6), respectively. The sequence of the tomato E4 promoter has also been published (Cordes, *et al.*, 1989), and the DNA sequences corresponding to segments pertinent to the invention are similarly provided in Figs. 2A and 2B (SEQ ID NO:1) and Fig. 10 (SEQ ID NO:6), respectively.

Polynucleotide segments used to construct the hybrid promoter can be obtained by PCR  
30 amplification of tomato genomic DNA, using primers designed on the basis of the information presented herein, coupled with a multi-functional sequence analysis program, *e.g.*, OLIGO version 5.0 for MacIntosh from National Biosciences, Inc. (Plymouth, MN), as detailed in Example 3.

Representative methods for isolating and characterizing an E4 and/or E8 promoter which can be applied for the purpose of obtaining a promoter segment of the present invention are described in  
35 International PCT Publication WO 95/35387.

The polynucleotide segments which make up the hybrid promoter were determined on the basis of expression results for transgenes driven by two representative hybrid promoters (Fig. 11), the long E8-E4 hybrid promoter (SEQ ID NO:1) and the short E8-E4 hybrid promoter (SEQ ID NO:6). Both versions of the hybrid promoter were highly effective in driving expression of an exemplary transgene coding for SAMase, as indicated by Western blot results shown in Figs. 6 and 7, which depict the results of studies on expression of SAMase driven by hybrid E4/E8 promoters derived from tomato in musk melon. The hybrid promoters were significantly more active in driving expression than either the tomato E8 or tomato E4 promoter.

The exemplary long E8/E4 hybrid promoter contains an E8 polynucleotide segment corresponding to nucleotides from about -2257 to -1103 of a tomato E8 promoter, while the short E8/E4 hybrid promoter contains an E8 polynucleotide segment corresponding to nucleotides from about -1529 to -847 of the E8 promoter. In expression experiments carried out in support of the invention, the short E8 promoter segment was generally found to function as effectively as the long E8 promoter segment to enhance overall activity of the hybrid promoter. This was surprising in view of previous reports indicating that DNA sequences necessary for both ethylene responsiveness and overall mRNA levels reside in the E8 promoter region encompassed by the longer E8/E4 hybrid promoter and not the shorter version (Deikman, *et al.*, 1992).

The E8 polynucleotide region encompassed by both versions of the hybrid promoter spans positions from about -2257 to -847 of the tomato E8 promoter. On the basis of these results, hybrid promoters of the invention will contain an E8 promoter-derived polynucleotide segment which preferably includes at least 30 contiguous nucleotides selected from this region. In one particular embodiment of the invention, the segment will include at least about 40 nucleotides selected from this region. In yet another embodiment, that the segment will include at least about 50 contiguous nucleotides. It will of course be appreciated that the E8-derived polynucleotide may contain fewer than 30 contiguous nucleotides, and in some instances, may contain from about 15-25 contiguous nucleotides.

The suitability of a particular E8 segment for use in constructs employing the hybrid promoter can be evaluated in expression experiments employing a heterologous reporter gene. A particular E8 segment selected according to the above guidelines is ligated to a downstream E4 promoter segment using the methods described herein. Expression levels of a suitable reporter gene driven by the resulting hybrid E8/E4 promoter are then compared to expression levels for the same gene regulated by the corresponding E8 or E4 promoter alone.

An E8 polynucleotide segment suitable for forming a hybrid promoter is one which, when combined with an E4 promoter polynucleotide segment corresponding to those described herein and placed in a hybrid promoter, drives expression of a reporter gene at a level of at least about 75-300%

of the expression level obtained using either an unmodified E4 or E8 gene promoter operably linked to said reporter.

A reporter gene, such as GUS ( $\beta$ -glucuronidase), can be used for this purpose. Expression of GUS protein can be easily measured by fluorometric, spectrophotometric or histochemical assays (Jefferson, 1987).

Turning now to the E4 polynucleotide segment of the hybrid promoter, an examination of the long E8/E4 hybrid promoter reveals an E4 polynucleotide segment corresponding essentially to the full-length E4 promoter, while the short E8/E4 hybrid promoter contains an E4 segment spanning from nucleotide positions -315 to +16 of the E4 promoter. As discussed above, both illustrative versions of the hybrid E8/E4 promoter were effective in directing expression of a heterologous gene (e.g., Examples 5 and 6). The high activity of the short E8/E4 hybrid was surprising, since studies on the E4 promoter alone indicate that both upstream and downstream elements are required for ethylene-responsive transcription (Xu, *et al.*, 1996).

The E4 polynucleotide region encompassed by both versions of the hybrid promoter spans nucleotide positions from about -1150 to +16 of the tomato E4 promoter. On the basis of these results, hybrid promoters of the invention will contain an E4 promoter-derived polynucleotide segment which preferably includes at least 200 contiguous nucleotides selected from this region. In one particular embodiment of the invention, the segment will include at least about 220 contiguous nucleotides selected from this region. In yet another embodiment, it is contemplated that the segment will include at least about 275 contiguous nucleotides. It will of course be appreciated that the E4-derived polynucleotide may contain fewer than 200 contiguous nucleotides, and may in some instances, contain from about 150-200 contiguous nucleotides.

Segments derived from the E4 gene will be selected and evaluated essentially as described above for the E8 component. Moreover, it will be appreciated that the above-described segments refer to functional equivalents thereof, and encompass promoters or promoter segments derived by means of ligation with operator regions, random or controlled mutagenesis, addition or duplication of enhancer sequences, addition or modification with synthetic linkers, and the like, having promoter activity similar to the E8-E4 hybrid promoter described herein or pertinent regions thereof.

#### IV. Chimeric Genes, Vector Construction and Plant Transformation

The E8/E4 hybrid promoter of the invention can be used to regulate expression of heterologous genes.

In support of the present invention, two exemplary chimeric genes containing an E8/E4 hybrid promoter sequence operably linked to a heterologous DNA sequence, were constructed, long E8/E4:SAMase (pAG-7162) and short E8/E4:SAMase (Examples 1 and 3). SAMase has been

predicted to function more efficiently if expressed (i) in high levels and (ii) in a tissue specific manner. Accordingly, the hybrid promoter described herein represents an ideal promoter to satisfy this objective, and can be used to express any heterologous gene fitting the above-description.

5           A.     Plant Transformation Vectors

Plant transformation vectors, containing an E8/E4 hybrid promoter/transcription-regulatory sequence, are constructed according to methods known in the art (see, for example, Houck and Pear, 1990, and Becker, *et al.*, 1992).

10           The present invention provides vectors suitable for the transformation of plants. The vectors, chimeric genes and DNA constructs of the present invention are also useful for the expression of heterologous genes. Transgenic plants, and their fruit products, carrying the chimeric genes of the present invention, may be a useful source of recombinantly-expressed material.

In one embodiment, the chimeric genes of the present invention have two components: (i) a hybrid E8/E4 promoter and (ii) a heterologous DNA coding sequence.

15           The vectors of the present invention may be constructed to carry an expression cassette containing an insertion site for DNA coding sequences of interest. The transcription of such inserted DNA is then under the control of a suitable E8/E4 hybrid promoter (*e.g.*, corresponding to SEQ ID NOs:1, 6).

20           Such expression cassettes may have single or multiple transcription termination signals at the coding-3'-end of the DNA sequence being expressed. The expression cassette may also include, for example, DNA sequences encoding (i) a leader sequence (*e.g.*, to allow secretion or vacuolar targeting), and (ii) translation termination signals.

25           Further, the vectors of the present invention may include selectable markers for use in plant cells (such as the *nptII* kanamycin resistance gene). The vectors may also include sequences that allow their selection and propagation in a secondary host, such as, sequences containing an origin of replication and a selectable marker. Typical secondary hosts include bacteria and yeast. In one embodiment, the secondary host is *Escherichia coli*, the origin of replication is a *colE1*-type, and the selectable marker is a gene encoding ampicillin resistance. Such sequences are well known in the art and are commercially available (*e.g.*, Clontech, Palo Alto, CA; Stratagene, La Jolla, CA).

30           The vectors of the present invention may also be modified to intermediate plant transformation plasmids that contain a region of homology to an *Agrobacterium tumefaciens* vector, a T-DNA border region from *Agrobacterium tumefaciens*, and chimeric genes or expression cassettes. Further, the vectors of the invention may comprise a disarmed plant tumor inducing plasmid of *Agrobacterium tumefaciens*. Other suitable vectors may be constructed using the promoters of the present invention  
35           and standard plant transformation vectors, which are available both commercially (Clontech, Palo

Alto, CA) and from academic sources (Waksman Institute, Rutgers, The State University of New Jersey, Piscataway, NJ).

The vectors of the present invention are useful for tissue and/or stage-specific expression of nucleic acid coding sequences in plant cells. For example, a selected peptide or polypeptide coding sequence can be inserted in an expression cassette of a vector of the present invention. The vector is then transformed into host cells, the host cells cultured under conditions to allow the expression of the protein coding sequences, and the expressed peptide or polypeptide isolated from the cells. Transformed progenitor cells can also be used to produce transgenic plants bearing fruit.

Further, the invention includes a method for producing a transgenic fruit-bearing plant, where fruit produced by the plant has a modified phenotype. In this method a chimeric gene is introduced (*e.g.*, by transformation) into progenitor cells of the plant. An exemplary chimeric gene is composed of (i) a DNA sequence encoding a gene product effective to modify a phenotypic characteristic of the plant, *e.g.*, to reduce ethylene biosynthesis in fruit produced by the plant, operably linked to (ii) a promoter whose expression is inducible, *e.g.*, during fruit ripening, by a plant cytokine, or by ethylene synthesis by the fruit. As above, the DNA sequence is heterologous to the promoter and the chimeric gene contains the appropriate regulatory elements necessary for expression in a plant. Transformed progenitor are grown cells to produce a transgenic plant bearing fruit. The method further includes transforming progenitor cells of the plant with a selectable vector containing the chimeric gene. The DNA sequences and promoters may be as described above.

In one aspect of the invention, fruit produced by such transgenic plants has a reduced level of ethylene synthesis by the fruit. The fruit then demonstrates a modified ripening phenotype in which the time course of ripening is delayed.

The vectors, chimeric genes and DNA constructs of the present invention can be sold individually or in kits for use in plant cell transformation and the subsequent generation of transgenic plants.

#### B. Isolation of E4 and E8 Promoters

The E4 and/or E8 promoter may be obtained from a homologue of the tomato E4 or E8 gene. To detect the presence of an E4 or E8 gene in various plant species, *e.g.*, strawberry, melon, carnation, cauliflower or raspberry, a southern blot experiment is carried out.

E4 or E8 homologues are identified in a Southern blot of the genomic DNA of a plant of interest, probed with a labeled DNA fragment containing the coding sequence of, *e.g.*, the tomato E4 or E8 gene.

Preferably, the probe is selected to contain the coding sequence of the tomato E4 or E8 gene rather than the promoter sequence, since coding sequences are typically more conserved from species to species than are promoter sequences.

Probes are generated from tomato genomic DNA using primer-specific amplification (Mullis, 1987; Mullis, *et al.*, 1987). The oligonucleotide primers are selected such that the amplified region includes the entire coding sequence of the tomato E4 or E8 gene. Primers may also be selected to amplify only a selected region of the E4 or E8 gene.

A promotor may also be isolated by (i) selecting first and second oligonucleotide primers corresponding to an upstream and a downstream region, respectively, of an E4 or E8 gene, (ii) amplifying a region of the E4 or E8 gene DNA between the first and second primers to generate probe molecules, (iii) contacting the probe molecules with a plurality of target DNA molecules derived from the genome of a selected fruit-bearing plant under conditions favoring specific hybridization between the probe molecule and a target molecule homologous to the probe molecule.

Alternatively, a probe can be made by isolating restriction-digest fragments containing the sequence of interest from plasmid DNA.

Such a probe is labeled with a detectable moiety to enable subsequent identification of homologous target molecules. Exemplary labeling moieties include radioactive nucleotides, such as <sup>32</sup>P-labeled nucleotides, digoxigenin-labeled nucleotides, biotinylated nucleotides, and the like, available from commercial sources.

In the case of a primer-amplified probe, labeled nucleotides may be directly incorporated into the probe during the amplification process. Probe molecules derived from DNA that has already been isolated, such as restriction-digest fragments from plasmid DNA, are typically end-labeled (Ausubel, *et al.*, 1992).

Target molecules, such as *Hind*III DNA fragments from the genomes of the above-listed plants, are electrophoresed on a gel, blotted, and immobilized onto a nylon or nitrocellulose filter. Labeled probe molecules are then contacted with the target molecules under conditions favoring specific hybridization between the probe molecules and target molecules homologous to the probe molecules, thereby identifying a target molecule having a DNA sequence homologous (having essentially the same sequence identity) to the tomato E4 or E8 gene, and isolating promoter sequences associated with the target molecule

Conditions favoring specific hybridization are referred to as moderately-to-highly stringent, and are affected primarily by the salt concentration and temperature of the wash buffer (Ausubel, *et al.*, 1992; Sambrook, *et al.*, 1989). Hybridization conditions are typically classified as moderately stringent, due to the low salt concentration, and are expected to preserve only specific hybridization interactions, allowing the identification and isolation of homologous genes in different plant species.

Following contacting, hybridization, and washing, target molecules with sequences substantially identical to the probe are identified by detecting the label on the probe. The label may be detected directly, for example, as in a radioactive label detected on autoradiograms, or it may be detected with a secondary moiety, for example, fluorescently-labeled streptavidin binding to a biotinylated probe.

Following the identification of plants containing E4 and/or E8 genes, the DNA encoding the genes, including the 5' regulatory regions, may be isolated from the respective species, by screening a genomic DNA library, *e.g.*, a library derived from a fruit-bearing plant.

The library of interest is screened with a probe containing sequences corresponding to the coding sequence of a known E4 gene, such as the tomato E4 gene. The screening is done using known methods (Ausubel, *et al.*, 1992; Sambrook, *et al.*, 1989).

Positive plaques or colonies are isolated, and the insert DNA is sequenced and compared to known E4 sequences. Clones containing inserts with sequences corresponding to genes with substantial sequence identity to tomato E4 are identified and, if necessary, used to obtain additional clones until the promoter region of interest is identified and further isolated. A DNA fragment containing an E4 or E8 promoter, or partial sequence thereof, may be isolated by digesting one of the lambda clones with selected restriction enzymes, *e.g.*, *HindIII* and *SacI*. This results in the generation of various fragments which may be purified by electrophoresis on agarose gels. The purified fragments may then be used to obtain the full-length promoter of the target E4 or E8 gene by hybridization using routine techniques known to those of skill in the art.

### C. Heterologous Genes

The methods and results described herein demonstrate the ability to provide high level tissue and/or stage specific gene expression in transgenic plants, where expression is regulated by a hybrid E8/E4 hybrid promoter. The E8/E4 hybrid promoter of the present invention includes a region or regions of DNA that regulates transcription of the immediately adjacent (downstream) gene to a specific plant tissue. According to methods of the present invention, heterologous genes are linked to the promoters of the present invention.

Exemplary heterologous genes for the transformation of plants include genes whose products are effective to reduce ethylene biosynthesis in specific tissues of those plants, *e.g.* the fruits. Some of these genes, including AdoMetase, are discussed above, and include S-adenosylmethionine hydrolase, aminocyclopropane-1-carboxylic acid (ACC) deaminase, ACC oxidase antisense molecule, ACC synthase antisense molecule, ACC oxidase cosuppression molecule, ACC synthase cosuppression molecule.

Other genes of interest that could be used in conjunction with the hybrid E8/E4 promoter include, but are not limited to other ripening modification genes in addition to AdoMetase. Representative examples of such genes include aminocyclopropane-1-carboxylic acid (ACC) deaminase (Klee, *et al.*, 1991; Sheehy, *et al.*, 1991), which degrades precursors of ethylene biosynthesis.

- 5 Ripening modification can also be achieved through the use of gene inactivation methods. Such methods may employ antisense or cosuppression-effecting genes of the ethylene biosynthetic pathway, *e.g.*, ACC synthase (Sato and Theologis, 1989) and ACC oxidase (Hamilton, *et al.*, 1990).

- Additional genes for use in the DNA constructs of the present invention include genes involved in conferring fungal resistance, *e.g.*, the polygalacturonase inhibiting protein, PGIP, from  
10 *Phaseolus vulgaris* (Toubart, *et al.*, 1992). Also contemplated are the use of modified forms of plant glucanase, chitinase and other pathogenesis related (PR) genes (Melchers, *et al.*, 1993, 1994; Ponstein, *et al.*, 1994; Woloshuk, *et al.*, 1991). Genes involved in conferring insect resistance may also be incorporated into the DNA constructs of the present invention. The expression of these products would be improved when used with a high-level, fruit-specific promoter such as the hybrid  
15 promoter of the present invention.

- In addition, antisense or cosuppression genes encoding proteins responsible for degradative processes in the fruit may also be used in conjunction with the promoters of the present invention. Examples of genes of this type include polygalacturonase, cellulase, and pectin methyl esterase (Schuch, 1994). In this way, inhibition of the specific degradation process is targeted only to  
20 ripening fruit.

- Other gene products which may be useful to express using the hybrid promoters of the present invention include gene products that are effective to modulate; (i) flowering, (ii) flavor (*e.g.*, thaumatin; GENBANK) or color modification (*e.g.*, products that modify lycopene synthesis, for example, arabidopsis lycopene cyclase; GENBANK), (iii) enzymes or other catalytic products such  
25 as ribozymes or catalytic antibodies that modify plant cell processes, (iv) ethylene production, such as antisense molecules, enzymes that degrade precursors of ethylene biosynthesis, catalytic products or cosuppression molecules, (v) fungal control, *e.g.*, alternative fungal control genes, (vi) production or levels of plant hormones, (vii) the cell cycle or cell division, and (viii) sucrose accumulation, such as the sucrose phosphate synthase gene (GENBANK) from corn.

- 30 Further, it is useful to modulate constitutive expression of some genes in specific tissues, such as expression of any gene that would be deleterious to the fruit if it were expressed constitutively, *e.g.*, genes which encode degradative enzymes that deplete necessary metabolites. Derivatives of the E4/E8 promoter can be used as on/off switches for the tissue and/or stage-specific expression of genes whose expression is under their control.

#### D. Methods of Transforming Plants

A number of methods, in addition to *Agrobacterium*-based methods, may be employed to elicit transformation of plant progenitor cells, such as electroporation, microinjection, and microprojectile bombardment. These methods are well known in the art (Comai and Coning, 1993; Klein, *et al.*, 1988; Miki, *et al.* 1987; Bellini, *et al.*, 1989) and provide the means to introduce selected DNA into plant genomes. Such DNA may include a DNA cassette which consists of a E8/E4 hybrid promoter functionally adjacent to heterologous sequences encoding a desired product, for example, AdoMetase coding sequences.

Representative methods for transforming tomato are described in International PCT Publication WO 95/35387. Representative methods for transforming strawberry or raspberry are described in International PCT Publication WO95/35388. Conventional protocols for transforming *Cucumis sp.* are described in Fang and Grumet, 1990; Valles and Lasa, 1994; Dong, *et al.*, 1991; Gonsalves, *et al.*, 1994; Yoshioka, *et al.*, 1992; Ayub, *et al.*, 1996.

Transformants and resulting transgenic cells and transgenic plants are identified and evaluated by standard methods (Mathews, *et al.*, 1995).

#### E. Expression in Heterologous Plant Systems

Experiments performed in support of the present invention demonstrate the versatility of the chimeric gene constructs of the invention (Examples 6, 7). The vector constructs of the present invention can be used for transformation and high level expression of heterologous sequences in transgenic plants. Further, the expression mediated by the promoters described herein appears to be tissue and/or stage-specific even in heterologous plants.

In looking now at experiments carried out in support of the invention, an evaluation of different promoters was conducted. Illustrative plant transformation experiments were carried out in muskmelon (*Cucumis melo*), using SAMase as the exemplary heterologous gene.

Transgenic melons were generated using *Agrobacterium*-mediated transformation and binary vectors containing a series of fruit and ripening -specific promoters from tomato (Table 2). Exemplary synthetic hybrid promoters containing different fruit-specific and ethylene-responsive promoter domains were prepared (*i.e.*, the long and short E8/E4 hybrid promoters) to determine their ability to enhance fruit and ripening specific gene expression.

Transgenic melons containing the SAMase gene were identified using PCR. Several transgenic events for each of the binary vectors were analyzed using Western blotting techniques to determine expression of the SAMase gene in ripening melon fruit.

Figures 6 and 7 indicate that neither the E8 nor the E4 promoters alone (Fig. 6) are equivalent to the E8/E4 hybrid promoters, in particular the long E8/E4 hybrid (Fig. 7), in their ability to drive

expression of SAMase in ripening melon fruit. While expression in the E4 and E8 driven constructs was observed to be weak-to-moderate, expression levels for the corresponding hybrid promoters was significantly greater, as determined by the intensity of the 17 kd AdoMetase band on the blot.

To further confirm and compare expression of the introduced transgenes, representative  
5 ethylene biosynthesis profiles were determined over a 4 day period for melon samples transformed with each of the representative constructs presented in Table 2.

Ethylene synthesis from the three different events (pAG-7142, pAG-7152, pAG-7162) are shown in Fig. 8, where entries followed by a (-) designation represent negative controls. In looking now at the results presented in Fig. 8, the pAG-7162-derived event (long E8/E4 hybrid promoter) is  
10 clearly reduced in its ability to produce ethylene during ripening, to an extent significantly greater than that of either of the E4 or E8-promoter driven events. Reduced ethylene synthesis and delayed ripening correlated with SAMase gene expression levels determined by Western blotting.

The long E8/E4 hybrid promoter-driven events demonstrate reduced ethylene biosynthesis, when compared to both the negative controls and to the other non-hybrid promoter-driven events.  
15 This is an indication of the greater expression activity of the hybrid promoter of the invention when compared to various non-hybrid promoters derived from different types of plant genes.

The use of the hybrid E4/E8 promoters cannot be considered limited to melon and tomato. The constructs and methods of the present invention are applicable to all higher plants, particularly fruits.

As demonstrated herein, the E4/E8 hybrid promoter sequences may be isolated from a type  
20 of plant other than the plant to be transformed. This is exemplified by the activity of an E4/E8 hybrid promoter composed of tomato-derived sequences which is effective to express a heterologous gene, *e.g.*, the SAMase gene in muskmelon. Alternatively, the E4/E8 hybrid promoter sequences may be isolated from the same type of plant as that which is transformed by a vector which contains an E4/E8  
25 hybrid promoter and a heterologous coding sequence, *e.g.* the SAMase gene. For example, a raspberry E4/E8 hybrid promoter may be operably linked to a heterologous gene, such as the SAMase gene, and used to transform raspberries.

The following examples illustrate, but in no way are intended to limit the scope of the present  
30 invention.

## Materials and Methods

### DNA Plasmids and *Agrobacterium* Binary Vector Constructions

Biological reagents were typically obtained from the following vendors: 5' to 3' Prime,  
5 Boulder, CO; New England Biolabs, Beverly, MA; Gibco/BRL, Gaithersburg, MD;  
Promega, Madison, WI; Clontech, Palo Alto, CA; and Operon, Alameda, CA.

Standard recombinant DNA techniques were employed in all constructions (Adams and Yang,  
1977; Ausubel, *et al.*, 1992; Hooykaas and Schilperoot 1985; Sambrook, *et al.*, 1989.

### Example 1

#### Long E8/E4 Hybrid Promoter (2.8kb) and Preparation of Intermediate Vector pAG-1762

##### A. Construction of intermediate vector pAG-1762

15 An intermediate vector, pAG-1762, which assembled all of the necessary elements into a  
moveable cassette was prepared as follows.

To obtain a portion of the tomato E8 promoter for use in preparing a hybrid promoter, a  
plasmid containing the 2.0 kb tomato E8 promoter, pAG-1742, was digested with *Xba*I and *Bam*HI  
using standard molecular biology protocols (Sambrook, *et al.*, 1989). The sequence of the 5' flanking  
region of the tomato E8 gene corresponding to 1 to -1098 and -2181 to -1098 has been previously  
20 described by Deikman, *et al.*, 1988 and Deikman, *et al.*, 1992, respectively.

A lambda genomic library from tomato was screened by standard methods with a tomato E8  
gene probe based on sequences described by Diekman *et al.* A lambda clone which hybridized to the  
probe was identified, plaque purified and sequenced. The E8 genomic clone was used as a source of  
the *Hind*III fragment that is the approximately -2257 to -1103 bp upstream region of the hybrid E8/E4  
25 promoter of the present invention, corresponding to nucleotides 1-1155 of SEQ ID NO:7. This  
fragment was inserted 5' of the approximately 1122 bp E8 promoter in pAG-5321 at the *Hind*III and  
*Xba*I sites (Figure 4).

The DNA sequence of the -2257 to -1103 base pair region is presented herein as nucleotides  
1 to 1160 of Figs. 2A and 2B. Following digestion, the excised tomato E8 promoter fragment was  
30 purified by electro-elution after agarose gel electrophoresis. The purified fragment was determined  
to contain the upstream sequence of the E8 promoter corresponding to nucleotides -2257 to -1098.  
The sequence corresponding to nucleotides -2257 to -1098 of the tomato E8 promoter is presented as  
nucleotides 1 to 1160 of SEQ ID NO:1 and 7.

To isolate a full-length E4 promoter for use in constructing a hybrid promoter, a 10.6 kb  
35 fragment containing the tomato E4 promoter was excised from a second plasmid, pAG-1752, by

treatment with *Xba*I and *Bam*HI. The sequence of the tomato E4 promoter has been published (Cordes, *et al.*, 1989), and the DNA sequence of the minus 1150 to plus 16 base pair region is presented as nucleotides 271 to 1437 of SEQ ID NO:8, which encompasses the full E4 gene.

5 The excised 10.6 kb fragment contained a tomato E4 promoter::SAMase construct. The sequence of the SAMase gene is described in U.S. Patent No. 5,589,623. The cloning and isolation of the SAMase gene is described in International Publication No. WO 95/35387. Also described therein are representative methods for isolating and characterizing a tomato E4 and/or E8 promoter, and details for preparing vector constructs of the type employed herein. The excised fragment, which contained the necessary sequences for bacterial selection and replication, was purified on agarose gel.

10 These two fragments were combined following the ligation protocol of Gibco/BRL to produce plasmid pAG-1762. This plasmid is shown schematically in Fig. 1.

### Example 2

#### Preparation of Binary Vector, pAG-7162 Containing Long E8/E4 Hybrid Promoter

15

The promoter/gene cassette from the intermediate vector, pAG-1762, was transferred into a binary vector as follows.

Plasmid pAG-1762 (Example 1) was digested with *Hind*III and *Kpn*I, resulting in a 2.8 kb fragment containing the long E8/E4 promoter (Figs. 2A, 2B; SEQ ID No:1) and SAMase. The excised fragment was gel purified by electrophoresis on agarose.

20

A second plasmid, pAG-7142 (Fig. 12) was digested with *Hind*III and *Kpn*I to provide a 12 kb fragment containing a terminator sequence and a selectable marker cassette (*i.e.*, the *np*tII kanamycin resistance gene driven by the raspberry E4 promoter) as well as the necessary sequences for bacterial selection and replication. The isolation, characterization, and sequence of the raspberry E4 (RE4) promoter, along with illustrative constructs describing the use of this promoter, are described in International PCT Publication WO 95/35388. The 12 kb fragment was purified by electrophoresis on agarose.

25

The purified 2.8 kb fragment containing the long E8/E4 promoter::SAMase construct was then ligated to the above-described purified 12 kb fragment to form plasmid pAG-7162, as illustrated schematically in Fig. 3.

30

**Example 3****Preparation of a Short E8/E4 Hybrid Promoter and an Intermediate Transfer Vector, pAG-126**

An intermediate vector combining truncated polynucleotide segments derived from the tomato E4 and E8 gene promoters to form a short E8/E4 hybrid promoter fused to the coding sequence for SAMase, was prepared as follows.

A tomato E4 promoter segment corresponding to nucleotides -315 to +16 was isolated from tomato DNA using PCR. In a similar fashion, PCR primers were designed to amplify a segment of the E8 promoter from -1529 to -847. The oligonucleotide primers were designed to incorporate restriction endonuclease sites at the ends of the amplified promoter fragments, for ready subcloning. The design of the primers was based on the published sequences of the tomato E8 and E4 genes as described above, using OLIGO version 5.0 for MacIntosh, a multi-functional sequence analysis program from National Biosciences, Inc. (Plymouth, MN).

**Table 1**

Primer	Sequence	ID No.
E4promod5	5'- <u>ACT CTT CCA</u> CAC TTT TCC CTC TA-3' underline= <i>Eco</i> I site	SEQ ID No: 2
E4promod3	5'- <u>CCC ATG GCT</u> CAA TCT CTA AT-3' underline= <i>Nco</i> I site	SEQ ID No: 3
E8promod5	5'- <u>CAA GCT TAA</u> AAT GTA CGA TGA GAG-3' underline= <i>Hind</i> III site	SEQ ID No: 4
E8promodLo	5'-AGT GTC <u>GAA GAG</u> GAC TAA ACC CGA AAA T-3' underline= <i>Eco</i> I site	SEQ ID No: 5

PCR was performed according to the manufacturer's procedure using Amplitaq (Perkin Elmer, Applied Biosystems Division, Foster City, CA), PCR buffer, tomato genomic DNA, and either pair of the above primers, employing the following (hot start) conditions:

- 1 cycle of 97°C for 5 minutes following addition of Amplitaq;
- 2 cycles of 97°C for 1 minute, 52°C for 1 minute and 72°C for 1 minute;
- 25 cycles of 94°C for 1 minute, 52°C for 1 minute and 72°C for 1 minute;
- 1 cycle of 72°C for 5 minutes; followed by cooling to 5°C.

The PCR-amplified fragments were each digested separately with *EaeI*. These two amplified fragments were then combined in a ligation reaction following conventional protocol (Gibco/BRL) utilizing ligase and buffer obtained from Gibco/BRL. The ligase reaction products were purified and the recovered DNA was subjected to digestion with restriction enzymes *NcoI* and *HindIII* (New England Biolabs, Beverly, MA).

The resulting short E8/E4 promoter fragment was then purified and ligated into a suitable plasmid vector. The vector, which contained the SAMase gene, was digested with *HindIII* and *NcoI* in order to orient the hybrid promoter immediately upstream of the SAMase gene, with both the promoter and gene positioned in the same 5' to 3' direction.

The resulting intermediate plasmid containing a short E8/E hybrid promoter::SAMase construct was designated pAG-126 and is presented in Fig. 4.

#### Example 4

##### Preparation of a Binary Vector Containing a Short E8/E4 Hybrid Promoter

The above-described promoter/gene cassette was transferred into a binary vector as follows.

Plasmid pAG-126 was digested with *HindIII* and *KpnI* to produce a 1.5 kb fragment containing the E8/E4 hybrid promoter coupled to the SAMase gene. The excised fragment was gel purified.

A binary plasmid vector, pAG-7142, containing a selectable marker gene cassette and a terminator sequence oriented downstream of the SAMase gene, was similarly treated with *HindIII* and *KpnI*. A schematic representation of vector pAG-7142 is presented as Fig. 12.

The 1.5 kb short E8/E4 hybrid promoter::SAMase fragment was then ligated to the binary vector to produce plasmid, pAG-7182, as shown in Fig. 5.

#### Example 5

##### A. Construction of Binary Vectors Containing Known Promoters

Binary vectors pAG-7142 (E4::SAMase) and pAG-7152 (E8::SAMase) were constructed following conventional protocols. Circular restriction maps of these plasmid vectors are shown in Figs. 12 and 9, respectively. The elements contained in each of the vectors were identical, with the exception of the promoter driving expression of SAMase.

### B. Transformation of *Cucumis Melo* (Muskmelon)

Cotyledon explants of a commercial cantaloupe variety were transformed according to known methods (Fang and Grumet, 1990; Valles and Lasa, 1994; Dong, *et al.*, 1991; Gonsalves, *et al.*, 1994; Yoshioka, *et al.*, 1992; Ayub, *et al.*, 1996).

5 The disarmed *Agrobacterium* strain EHA105 (Hood, *et al.*, 1993) was used to introduce the above-described binary vectors into plants. The disarmed *Agrobacterium* strain was cocultivated with melon cotyledon tissue explants. Primary transformants were selected on the basis of their capacity to regenerate and develop roots on media containing the antibiotic, kanamycin.

10 The presence of the SAMase gene in the putative transgenic plants was confirmed using the polymerase chain reaction and primers specific for the SAMase gene.

Primary transgenic plants were grown in a greenhouse. Individual melon fruit were harvested at a stage when the fruit had initiated the process of abscission. Harvested fruit were stored under refrigeration for 7 days. Samples were then frozen at -70°C for storage prior to gene expression studies.

15

### Example 6

#### Western Blot Analysis of SAMase Gene Expression

Protein lysates were prepared from the melon pericarp fruit tissue of each sample and were free of any rind tissue. Using liquid nitrogen and a mortar and pestle, one gram tissue samples were  
20 ground to a fine powder, mixed with Laemmli SDS sample buffer (Laemmli, 1970) and heated at 95-100°C to denature the proteins.

Protein concentrations were determined by precipitating the proteins from the melon lysate with trichloroacetic acid, resolubilizing the proteins, followed by a quantification of protein concentration using Pierce's BCA (bicinchoninic acid) assay kit, according to the manufacturer's  
25 instructions. Sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS PAGE) was performed using Laemmli's denaturing (SDS) discontinuous gel electrophoresis method (Laemmli, 1970).

Proteins resolved by SDS PAGE were transblotted on to Millipore's Immobilon-P PVDF membrane in Tris-glycine/15% methanol buffer at 100 volts for 2 hours. The membranes were then treated with 5% (w/v) powdered milk in phosphate buffered saline (PBS) containing Tween 20 for 2  
30 hours at room temperature.

The primary antibody treatment was performed using 1 µg/ml of an anti-SAMase mouse monoclonal antibody in 3.5% powdered milk/PBS-Tween 20 for 2 hours at room temperature. The secondary antibody, a horseradish peroxidase-labeled goat anti-mouse IgG+IgM(H+L) polyclonal antibody (Kirkegaard and Perry Laboratories, Inc., Gaithersburg, MD), was used to treat the  
35 membrane at 0.1µg/ml in 3.5% powdered milk/PBS-Tween 20 for 2 hours at room temperature.

Membrane bound antibody was detected using NEN/DuPont's Renaissance chemiluminescence reagent in a 1 minute incubation as per the manufacturer's instructions. The blot was then exposed to Kodak BioMax MR X-ray film for at least one hour.

5

**Table 2**

Summary of *Agrobacterium* Vectors Used  
to Transform Melon

10

Construct ID#	Promoter::SAMase	Promoter::Selectable Marker
pAG-7142	Tomato E8::SAMase	RE4::nptII
pAG-7152	Tomato E4::SAMase	RE4::nptII
pAG-7162	Long E8/E4::SAMase	RE4::nptII
pAG-7182	Short E8/E4::SAMase	RE4::nptII

15

All of the binary vector constructs are identical with the exception of the promoter controlling expression of the exemplary SAMase gene.

Several transgenic events for each of the binary vectors were analyzed using Western blots to determine expression of the SAMase gene in ripening melon fruit.

20

Figs. 6 and 7 indicate that neither the E8 nor the E4 promoters alone are equivalent to the E8/E4 hybrid promoters, in particular the long E8/E4 hybrid, in their ability to drive expression of SAMase in ripening melon fruit.

### Example 7

25

#### Ethylene Production in Melon Transformants

Representative ethylene biosynthesis profiles were determined over a 4 day period. At day zero, transformed melon was at a physiological ripening stage known as ¼ slip. "Slip", in melon ripening terminology, refers to the abscission of the fruit from the peduncle of stem region which attaches the melon to the vine. Thus, ¼ slip signifies the initiation of the ripening process.

30

The assay for ethylene evolution in transgenic melon fruit is performed by placing a rubber cup over the stem end of individual fruit and sampling 0.25 ml aliquots for gas chromatographic analysis after approximately four hours (Ayub, *et al.*, 1996). A Hewlett Packard 6890 (Hewlett Packard, Palo Alto, CA) gas chromatograph with a flame ionization detector and a 6 ft Haysep-D column was used for ethylene measurements. This system, when utilized in combination with an HP Vectra computer and the current version of "CHEMSTATION" (Hewlett Packard) allows measurement of ethylene concentrations as low as 0.1 ppm.

35

Ethylene synthesis from the three different events (pAG-7142, pAG-7152, pAG-7162) are shown in Fig. 8, where entries followed by a (-) designation represent negative controls. The results are typical of other transgenic events that have been analyzed to date.

In looking now at the results presented in Fig. 8, the pAG-7162-derived event (long E8/E4  
5 hybrid promoter) is clearly reduced in its ability to produce ethylene during ripening, to an extent significantly greater than that of either of the E4 or E8-promoter driven events. The long E8/E4 hybrid promoter-driven events demonstrate reduced ethylene biosynthesis, when compared to both the negative controls and to the other non-hybrid promoter-driven events.

**IT IS CLAIMED:**

1. A DNA construct comprising, in a 5' to 3' direction:
  - (i) a hybrid promoter comprising a segment of at least 30 contiguous nucleotides selected from  
5 the region extending from nucleotides 1 to 1395 of SEQ ID NO:7 of the tomato E8 gene promoter  
or the functional equivalent thereof fused to a polynucleotide segment of at least 200 contiguous  
nucleotides selected from the region extending from nucleotides 271 to 1437 of SEQ ID NO:8, or the  
functional equivalent thereof and  
a heterologous DNA coding sequence operably linked to said hybrid promoter,  
10 whereby said hybrid promoter is effective to drive expression of a reporter gene at a level of  
at least about 75-300% of the expression level obtained using either an unmodified tomato E4 or  
tomato E8 gene promoter operably linked to said reporter.
2. The DNA construct of claim 1, wherein said tomato E8 promoter segment comprises  
15 nucleotides 729 to 1411 of SEQ ID NO:7.
3. The DNA construct of claim 1, wherein said tomato E4 promoter segment comprises  
nucleotides 1107 to 1437 of SEQ ID NO:8.
- 20 4. The DNA construct of claim 1, wherein said hybrid promoter comprises a tomato E8  
promoter segment consisting essentially of nucleotides 729 to 1411 of SEQ ID NO:7 and a tomato E4  
promoter segment consisting essentially of nucleotides 1107 to 1437 of SEQ ID NO:8.
5. The DNA construct of claim 1, wherein said tomato E8 promoter segment comprises  
25 nucleotides 1 to 1156 of SEQ ID NO:7.
6. The DNA construct of claim 1, wherein said tomato E4 promoter fragment comprises  
nucleotides 271 to 1437 of SEQ ID NO:8.
- 30 7. The DNA construct of claim 1, wherein said hybrid promoter comprises a tomato E8  
promoter segment consisting essentially of nucleotides 1 to 1156 of SEQ ID NO:7 and a tomato E4  
promoter segment consisting essentially of nucleotides 271 to 1437 of SEQ ID NO:8.

8. A DNA construct of claim 1, wherein said heterologous DNA sequence encodes a product effective to reduce ethylene biosynthesis or delay ripening when expressed in fruit from a fruit-bearing plant.

5           9. A DNA construct of claim 8, wherein said heterologous DNA sequence encodes a product selected from the group consisting of S-adenosylmethionine hydrolase (SAMase), phytoene synthase, aminocyclopropane-1-carboxylic acid (ACC) deaminase, ACC oxidase antisense molecule, ACC synthase antisense molecule, ACC oxidase cosuppression molecule, and ACC synthase cosuppression molecule.

10

10. A DNA construct of claim 1, wherein said heterologous DNA coding sequence corresponds to a pathogenesis-resistance gene.

11. A DNA construct of claim 1, wherein said hybrid promoter is an ethylene-responsive  
15 promoter.

12. A DNA construct of claim 1, wherein said hybrid promoter is a fruit-specific promoter.

13. A plant transformation vector containing the DNA construct of any of claims 1-12.

20

14. A dicotyledonous plant cell transformed with the DNA construct of any of claims 1-12.

15. A kit for use in plant transformation comprising the vector of claim 13.

25

16. A method for producing a transgenic plant, comprising the steps of:

(i) introducing into progenitor cells of said plant a DNA construct of any of claims 1-12 to produce transformed plant cells, and

(ii) growing the transformed progenitor plant cells to produce a transgenic plant capable of expressing said heterologous DNA coding sequence.

30

17. A method of claim 16, wherein (i) said transgenic plant is a fruit-bearing plant, (ii) said growing produces a transgenic plant bearing fruit, and (iii) said heterologous DNA coding sequence is preferentially expressed in fruit from said plant.

18. A method of claim 16, wherein said plant is selected from the group consisting of *Cucumis sp.*, raspberry, strawberry and tomato.

19. A method of claim 16, wherein said plant is *Cucumis sp.*

5

20. A method for inhibiting fruit ripening in fruit from a fruit-bearing plant comprising:

(i) introducing into progenitor cells of said plant a DNA construct of claim 8 to produce transformed plant cells,

(ii) growing the transformed progenitor plant cells to produce a transgenic plant bearing fruit,  
10 whereby said heterologous DNA sequence is expressed in said fruit to form a product effective to delay the timecourse of ripening of the fruit relative to non-transformed fruit.

21. The method of claim 20, wherein (i) the tomato E8 promoter fragment comprises nucleotides 1 to 1156 of SEQ ID NO:7, (ii) the tomato E4 promoter segment comprises nucleotides  
15 271 to 1437 of SEQ ID NO:8, and (iii) said heterologous DNA coding sequence encodes SAMase.

22. The method of claim 20, wherein (i) the tomato E8 promoter comprises nucleotides 712 to 1395 of SEQ ID NO:7, (ii) the tomato E4 promoter comprises nucleotides 1107 to 1437 of SEQ ID NO:8, and (iii) said heterologous DNA coding sequence encodes SAMase.

20


23. The method of claim 20, wherein said fruit is melon.

24. A method for conferring enhanced expression activity to a tomato E4 promoter which comprises:

25 fusing, in an upstream orientation to a tomato E4 promoter polynucleotide segment of at least 200 contiguous nucleotides selected from the region extending from nucleotide positions 271 to 1437 of SEQ ID NO:8, a polynucleotide segment of at least 30 contiguous nucleotides selected from the region extending from nucleotide positions 1 to 1395 of SEQ ID NO:7 of the tomato E8 gene promoter, to thereby form a hybrid E8-E4 promoter capable of regulating expression of a heterologous  
30 gene operably linked thereto,

whereby said hybrid promoter is effective to drive expression of said heterologous gene at a level of at least about 75-300% of the expression level obtained using an unmodified tomato E4 gene promoter operably linked to said heterologous gene.

1/13

**Fig. 1**

2/13

AAGCTTTAAT TGGTTGAGAT TGAACGTAAT TCAAATTATT CTGAGCCCAA ACCCTTAAAA TTCTAGGCGG  
TTATCTTTGT TTGAATTCAT TTTTGACATC CCTAATGATA TTGTTACAGT AATTAAGTTT TGTGGAAGTG  
AGAGAGTCCA ATTTTGATAA GAAAAGAGTC AGAAAACGTA ATATTTTAAA AGTCTAAATC TTTCTACAAA  
TAAGAGCAAA TTTATTTATT TTTTAATCCA ATAAATATTA ATGGAGGACA AATTCAATTC ACTTGGTTGT  
AAAATAAACT TAAACCAATA ACCAAAGAVC TAATAAATCT GAAGTGGAAT TATTAAGGAT AATGTACATA  
GACAATGAAG AAATAATAGG TTCGATGAAT TAATAATAAT TAAGGATGTT ACAATCATCA TGTGCCAAGT  
ATATACACAA TATTCTATGG GATTTATAAT TTCGTTACTT CACTTAACTT TTGCGTAAAT AAAACGAATT  
ATCTGATATT TTATAATAAA ACAGTTAATT AAGAACCATC ATTTTAAACA ACATAGATAT ATTATTTCTA  
ATAGTTTAAT GATACTTTTA AATCTTTTAA ATTTTATGTT TCTTTTAGAA AATAAAAATT CAAAAAATT  
AAATATATTT ACAAAAAC TAATCAAACA CAACCTCATA TATTAAAAGC AAAATATATT TTGAAAATTT  
CAAGTGTCCT AACAAATAAG ACAAGAGGAA AATGTACGAT GAGAGACATA AAGAGAACTA ATAATTGAGG  
AGTCCTATAA TATATAATAA AGTTTATTAG TAAACTTAAT TATTAAGGAC TCCTAAAATA TATGATAGGA  
GAAAATGAAT GGTGAGAGAT ATTGGAAGAA TTAATAATTA AGGATNTTAA AATATATGGT AAAAGATAGG  
CAAAGTATCC ATTATCCCCT TTAACTTGA AGTCTACCTA GCGCATGTG AAAGGTTGAT TTTTGTGAC  
GTCATATAGC TATAACGTAA AAAAAGAAAG TAAAATTTTT AATTTTTTTT AATATATGAC ATATTTTAAA  
CGAAATATAG GACAAAATGT AAATGAATAG TAAAGGAAAC AAAGATTAAT ACTTACTTTG TAAGAATTTA  
AGATAAATTT AAAATTTAAT AGATCAACTT TACGTAAAG TAACTTGGG TGGGTCAAGA CCCAACTCGA  
TTTCTGTTCA ACCCATTTTA ATATTTCTAT TTTCAACCTA ACCCGCTCAT TTGATACCCC TACAAATATC  
ATATTTGTGT GTGAAATATT TTTTGGGCTG GAGAGAGAGG CCCCAGGGG AGTGGAGGGG TGGGGTGGGG  
AGAGAGAGCG AGAAAGAGTG GAGAGAGAAA TTTGATATGA AATCCTACAT ATATTACAGA TTGTAATGTT


**Fig. 2A**

3/13


CTAAACTATA ACGATTTGTC ATAAACACAT ATCATGGATT TGTCTTTTGT TGTAATTTTC CCAATTGTAA  
ATAGGACTTC GTTATTTGAA ACTTGAAAGT GAAGTCACAT AGATTAAAGTA CAAACATTAA TTAAAGACCG  
TGGTGAATG ATAAATATTT ATTTATCTTT AATTAGTTAT TTTTTTGGGA GCTCTTTATT CCAATGTGAG  
ACTTTTGCGA CATATATTCA AATTTAATCG AATCACAATA TGTATTAGAT TGATAAAAAA ATAATTTTTT  
TACAATGTTA GTTGAGACTC ATAACCTACT GCCTATTGGT AATCTATGAC TCCTAATTCC TTAATTATTT  
AAATATATCA TCTTGATCGT TAACAAAGTA ATTTGAAAG ACCACGAGTA AGAAGACAAA CGAGAATACC  
AAAAAATTCA AAAATTTAAT GTGATTTGGT CAATCGATCT ACGTCCATAA AGGAGATGAG TAATCTACTA  
TAAATATGAG AGTACAAAAT ACAGAGAGAA ACAACCTCAA CTAATTCACT CGGAATACAT GAGAAGTTCA  
CACAAGTGAT AACGTATCAA ACTTGTGACC CACACTTTTC CCTCTAACCA AAGCTCTTAA AACTATATTG  
TGAATGCTGA TTAAGTTAAA CGAAACAGTC CTAAATCTTT TCCGTCCTAT GAGAAACAAG ATTAATCAAT  
TCACAATTTT TTTAAAAAGA AAAACCTGTA AGAAATTTAG GCAAACAAA CCTAACACAA GTTTGTTTTT  
GTTTTTACTA CCAACAAGAA ATTCAAATGG CAAATGTATA ACGCATCTTA GCTAATTATA TGACCAGATT  
CAGATTAATA TACATCTTCA CCCATGCAAT CCATTTCTAT ATAAAGAAAC ATACACGAAC TTGATATTAT  
TAGAGATTGA GCCATGG

**Fig. 2B**


4/13

**Fig. 3**


5/13

**Fig. 4**

6/13

**Fig. 5**

7/13



**Fig. 6**

8/13






Fig. 7

9/13

**Fig. 8**

10/13


**Fig. 9**

11/13

CTTAAATGTACGATGAGAGACATAAAGAGAACTAATAATTGAGGAGTCCTATAATATATAAT  
AAAGTTTATTAGTAAACTTAATTATTAAGGACTCCTAAAAATATATGATAGGAGAAAATGAATG  
GTGAGAGATATTGGAACCTTAATAATTAAGGATNTTAAAAATATATGGTAAAAGATAGGCAAA  
GTATCCATTATCCCCTTTTAACCTGAAGTCTACCTAGGCGCATGTGAAAGGTTGATTTTTTGT  
CACGTCATATAGCTATAACGTAAAAAAGAAAGTAAATTTTTAATTTTTTTAATATATGAC  
ATATTTTAAACGAAATATAGGACAAAATGTAAATGAATAGTAAAGGAAACAAAGATTAATACT  
TACTTTGTAAGAATTTAAGATAAATTTAAATTTAATAGATCAACTTTACGTCTAGAAAGACC  
CATATCTAGAAGGAATTCACGAAATCGGCCCTTATTCAAAAATAACTTTTAAATAATGAATT  
TTAAATTTAAGAAATAATATCCAATGAATAAATGACATGTAGCATTTTACCTAAATATTTCA  
ACTATTTAATCCAATATTAATTTGTTTTATTCCCAACAATAGAAAGTCTTGTGCAGACATTT  
AATCTGACTTTTCCAGTACTAAATATTAATTTCTGAAGATTTTCGGGTTTAGTCCTCTTCGA  
CACTTTTTTACAATTTTTTTAAAAAGAAAACCTGTAAGAAATTTAGGCAAAACAAAACCTAA  
CACAAGTTTGTTTTTGTTTTTTACTACCAACAAGAAATTCAAATGGCAAATGTATAACGCATCT  
TAGCTAATTATATGACCAGATTCAGATTAATATACATCTTCACCCATGCAATCCATTTCTATA  
TAAAGAAACATACACGAACCTTGATATTATTAGAGATTGAGCCATGG


**Fig. 10**

12/13



**Fig. 11**

13/13

**Fig. 12**

## SEQUENCE LISTING

&lt;110&gt; Agritope, Inc.

&lt;120&gt; Synthetic Hybrid Plant Promoters

&lt;130&gt; 4257-0016.41

&lt;140&gt; unassigned

&lt;141&gt; 1998-09-18

&lt;150&gt; 60/059,234

&lt;151&gt; 1997-09-18

&lt;160&gt; 10

&lt;170&gt; FastSEQ for Windows Version 3.0

&lt;210&gt; 1

&lt;211&gt; 2327

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; promoter

&lt;222&gt; (1) ... (2327)

&lt;223&gt; synthetic DNA promoter sequence

&lt;400&gt; 1

aagctttaat	tgggtgagat	tgaacgtaat	tcaaattatt	ctgagcccaa	acccttaaaa	60
ttctagcg	ttatcttgg	ttgaattcat	ttttgacatc	cctaatagata	ttgttcacgt	120
aattaagttt	tgtggaagt	agagagtgcca	attttgataa	gaaaagagtc	agaaaacgta	180
atattttaaa	agtctaaatc	tttctacaaa	taagagcaaa	tttattttatt	ttttaatcca	240
ataaatatta	atggaggaca	aattcaattc	acttggttgt	aaaataaaact	taaaccaata	300
accaaagavc	taataaatct	gaagtgggaat	tattaaggat	aatgtacata	gacaatgaag	360
aaataatagg	ttcgatgaat	taataataat	taaggatggt	acaatcatca	tgtgccaaagt	420
atatacacaa	tattctatgg	gatttataat	ttcgttactt	cacttaactt	ttgcgtaaat	480
aaaacgaatt	atctgatatt	ttataataaa	acagttaatt	aagaaccatc	atttttaaca	540
acatagatat	attatttcta	atagtttta	gatactttta	aatcttttaa	attttatggt	600
tcttttagaa	aataaaaaatt	caaaaaaatt	aaatatattt	acaaaaacta	caatcaaaca	660
caacttcata	tattaaaagc	aaaatatatt	ttgaaaattt	caagtgtcct	aacaaataag	720
acaagaggaa	aatgtacgat	gagagacata	aagagaacta	ataattgagg	agtcctataa	780
tatataataa	agttttattag	taaacttaat	tattaaggac	tcctaaaata	tatgatagga	840
gaaaatgaat	ggtgagagat	attggaaaac	ttaataatta	aggatnttaa	aatatatggt	900
aaaagatagg	caaagtatcc	attatcccct	tttaacttga	agtctaccta	ggcgcagtgt	960
aaaggttgat	tttttgtcac	gtcatatagc	tataacgtaa	aaaaagaaaag	taaaattttt	1020
aatttttttt	aatatatgac	atatttttaa	cgaaatatag	gacaaaatgt	aaatgaatag	1080
taaaggaaac	aaagattaat	acttactttg	taagaattta	agataaattt	aaaatttaat	1140
agatcaactt	tacgttaaag	taaacttggg	tgggtcaaga	cccaactcga	tttctgttca	1200
acccatttta	atatttctat	tttcaaccta	acccgctcat	ttgatacccc	tacaaatatc	1260
atatttgtgt	gtgaaatatt	ttttgggctg	gagagagagg	ccccgagggg	agtggagggg	1320
tgggggtggg	agagagagcg	agaaaagagt	gagagagaaa	tttgatatga	aatcctacat	1380
atattacaga	ttgtaatggt	ctaaactata	acgatttgct	ataaacacat	atcatggatt	1440
tgtctttttg	tgtaattttc	ccaattgtta	ataggacttc	gttatttgaa	acttgaaagt	1500
gaagtcacat	agattaagta	caaacattaa	ttaaagaccg	tgggtggaatg	ataaatattt	1560
atattatctt	aattagttat	ttttttggga	gctctttatt	ccaatgtgag	actttttgca	1620
catatatatt	caatttaacg	aatcacata	tgtattagat	tgataaaaaa	ataatttttt	1680
tacaatgtta	gttgagactc	ataacttact	gcctattggt	aatctatgac	tcctaattcc	1740
ttaattat	aaatatatca	tcttgatcgt	taacaaagta	atttcgaaag	accacgagta	1800

agaagacaaa	cgagaatacc	aaaaaattca	aaaattttaat	gtgatttggt	caatcgatct	1860
acgtccataa	aggagatgag	taatctacta	taaatatgag	agtacaaaat	acagagagaa	1920
acaacctcaa	ctaattcact	cggaatacat	gagaagttca	cacaagtgat	aacgtatcaa	1980
acttgtagcc	cacacttttc	cctctaacca	aagctcttaa	aactatattg	tgaatgctga	2040
ttaagttaaa	cgaaacagtc	ctaaatcttt	tccgtcctat	gagaaacaag	attaatcaat	2100
tcacaatttt	tttaaaaaaga	aaaacctgta	agaaatttag	gcaaacaaaa	cctaacacaa	2160
gtttgttttt	gtttttacta	ccaacaagaa	attcaaattg	caaattgtata	acgcatctta	2220
gctaattata	tgaccagatt	cagattaata	tacatcttca	cccatgcaat	ccatttctat	2280
ataaagaaac	atacacgaac	ttgatattat	tagagattga	gccatgg		2327

&lt;210&gt; 2

&lt;211&gt; 23

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (1)...(23)

&lt;223&gt; synthetic oligonucleotide

&lt;400&gt; 2

actcttccac acttttccct cta

23

&lt;210&gt; 3

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (1)...(20)

&lt;223&gt; synthetic oligonucleotide

&lt;400&gt; 3

cccatggctc aatctctaata

20

&lt;210&gt; 4

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (1)...(24)

&lt;223&gt; synthetic oligonucleotide

&lt;400&gt; 4

caagcttaaa atgtacgatg agag

24

&lt;210&gt; 5

&lt;211&gt; 28

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; primer\_bind

&lt;222&gt; (1)...(28)

&lt;223&gt; synthetic oligonucleotide

&lt;400&gt; 5

agtgctgaag aggactaaac ccgaaaaat

28

<210> 6  
 <211> 1028  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> promoter  
 <222> (1)...(1028)  
 <223> synthetic DNA promoter sequence

<400> 6

cttaaaatgt	acgatgagag	acataaagag	aactaataat	tgaggagtcc	tataatatat	60
aataaagttt	attagtaaac	ttaattatta	aggactccta	aaatatatga	taggagaaaa	120
tgaatggtga	gagatattgg	aaaacttaat	aattaaggat	nttaaaatat	atggtaaaag	180
ataggcaaag	tatccattat	ccccttttaa	cttgaagtct	acctaggcgc	atgtgaaagg	240
ttgatTTTTT	gtcacgtcat	atagctataa	cgtaaaaaaa	gaaagtaaaa	tttttaattt	300
tttttaatat	atgacatatt	ttaaacgaaa	tataggacaa	aatgtaaatg	aatagtaaag	360
gaaacaaaga	ttaatactta	ctttgtaaga	atthaagata	aatTTTaaat	ttaatagatc	420
aactttacgt	ctagaaagac	ccatatctag	aaggaatttc	acgaaatcgg	cccttatcca	480
aaaataactt	ttaaataatg	aattttaaat	tttaagaaat	aatatccaat	gaataaatga	540
catgtagcat	tttacctaaa	tatttcaact	attttaatcc	aatattaatt	tgtttttattc	600
ccaacaatag	aaagtcttgt	gcagacattt	aatctgactt	ttccagtact	aaatattaat	660
tttctgaaga	ttttcgggtt	tagtctctct	cgacactttt	ccctctaacc	aaagctctta	720
aaactatatt	gtgaatgctg	attaagttta	acgaaacagt	cctaaatcct	ttccgtccta	780
tgagaaacaa	gattaatcaa	ttcacaattt	ttttaaaaaa	aaaaacctgt	aagaaattta	840
ggcaaacaaa	acctaacaca	agtttgTTTT	tgtttttact	accaacaaga	aattcaaattg	900
gcaaatgtat	aacgcattct	agctaattat	atgaccagat	tcagattaat	atacatcttc	960
acccatgcaa	tccatttcta	tataaagaaa	catacacgaa	cttgatatta	ttagagattg	1020
agccatgg						1028

<210> 7  
 <211> 2298  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> synthetic hybrid DNA sequence

<400> 7

aagctttta	tggttgagat	tgaacgtaat	tcaaattatt	ctgagcccaa	acccttaaaa	60
ttctagggc	ttatctttgt	ttgaattcat	ttttgacatc	cctaattgata	ttgttcacgt	120
aattaagttt	tgtggaagt	agagagtcca	atTTtgataa	gaaaagagtc	agaaaacgta	180
atatttttaa	agtctaaatc	tttctacaaa	taagagcaaa	tttattttatt	ttttaatcca	240
ataaatatta	atggaggaca	aattcaattc	acttggttgt	aaaataaact	taaaccaata	300
accaaaganc	taataaatct	gaagtggaat	tattaaggat	aatgtacata	gacaatgaag	360
aaataatagg	ttcgatgaat	taataataat	taaggatgtt	acaatcatca	tgtgccaaagt	420
atatacacaa	tattctatgg	gattttataat	ttcgttactt	cacttaactt	ttgcgtaaat	480
aaaacgaatt	atctgatatt	ttataataaa	acagttaatt	aagaaccatc	atTTTTaaca	540
acatagatat	attattttcta	atagtttaat	gatactttta	aatcttttaa	atTTttatgtt	600
tcttttagaa	aataaaaaatt	caaaaaaatt	aaatatattt	acaaaaacta	caatcaaaca	660
caacttcata	tattaaaagc	aaaatatatt	ttgaaaattt	caagtgtcct	aacaaataag	720
acaagaggaa	aatgtacgat	gagagacata	aagagaacta	ataattgagg	agtcctataa	780
tatataataa	agttttattag	taaacttaat	tatttaaggac	tcctaaaaata	tatgatagga	840
gaaaatgaat	ggtgagagat	attggaaaac	ttaataatta	aggatnttaa	aatatatggt	900
aaaagatagg	caaagtatcc	attatcccct	tttaacttga	agtctaccta	ggcgcatgtg	960
aaaggttgat	tttttgtcac	gtcatatagc	tataacgtaa	aaaaagaaaag	taaaattttt	1020
aatTTTTttt	aatatatgac	atatttttaa	cgaatatatag	gacaaaatgt	aaatgaatag	1080
taaaggaaac	aaagattaat	acttactttg	taagaattta	agataaattt	aaaattttaat	1140
agatcaactt	tacgtctaga	aagaccata	tctagaagga	atTTTcacgaa	atcgccctt	1200
attcaaaaat	aactttttaa	taatgaattt	taaatttttaa	gaaataatat	ccaatgaata	1260

aatgacatgt	agcatttttac	ctaaatattt	caactatttt	aatccaatat	taatttgttt	1320
tattcccaac	aatagaaaagt	cttgtgcaga	catttaattct	gactttttcca	gtactaaata	1380
ttaattttct	gaagattttc	gggttttagtc	cacaagtttt	agtgagaagt	tttgctcaaa	1440
attttaggtg	agaaggtttg	atatttatct	tttgttaaat	taatttatct	aggtgactat	1500
tatttattta	agtagaaaat	catatcatta	cttttgccaa	cttgtagtca	taataggagt	1560
agggtgtatat	gatgaaggaa	taacaaggtt	cagtgaagt	attaaaataa	aatataattt	1620
agggtgtacat	caaataaaaa	ccttaaagtt	tagaaaggca	ccgaataatt	ttgcatagaa	1680
gatatttagta	aattttataaa	aataaaaagaa	atgtagttgt	caagttgtct	tctttttttt	1740
ggataaaaaat	agcagttggc	ttatgtcatt	cttttacaac	ctccatgcca	cttgtccaat	1800
tgttgacact	taactaatta	gtttgattca	tgtatgaata	ctaaataatt	ttttaggact	1860
gactcaaata	tttttatatt	atcatagtaa	tatttatcta	attttttagga	ccacttatta	1920
ctaaataata	aattaactac	tactatatta	ttgttgtgaa	acaacaacgt	tttggttgtt	1980
atgatgaaac	gtacactata	tcagtatgaa	aaattcaaaa	cgattagtat	aaattatatt	2040
gaaaatttga	tatttttcta	ttcttaatca	gacgtatttg	gtttcatatt	ttaaaaagg	2100
actaaactta	gaagagaagt	ttgtttgaaa	ctacttttct	ctctttcttg	ttcccatttc	2160
tctcttagat	ttcaaaaagt	gaactacttt	atctctttct	ttgttcacat	tttattttat	2220
tctattataa	atatggcatc	ctcatattga	gattttttaga	aattattcta	atcattcaca	2280
gtgcaaaaaga	ccatggaa					2298

&lt;210&gt; 8

&lt;211&gt; 2796

&lt;212&gt; DNA

&lt;213&gt; Solanum esculentum

&lt;400&gt; 8

gaattctcaa	ttgagcccaa	ttcaatctcc	aatttcaacc	cgttttaaaa	ctttttatta	60
agatatgttt	ctatatgtga	agtatgaatt	attatctatt	taacatcttt	taggatttat	120
ctatccattt	gctacttttt	taacaaaaaa	ttcttgagtg	aaaattcaaa	ttgtgattat	180
aaaagttaaa	tatcaatatg	ttaaattatt	aagattaatc	gggtcaaat	ggcgggtcaa	240
ggcccaattc	tttttttagcc	catttaagct	caaagtaaac	ttgggtgggt	caagacccaa	300
ctcgatttct	gttcaaccce	ttttaatatt	tctattttca	acctaaccg	ctcattttgat	360
acccctacaa	atatcatatt	tgtgtgtgaa	atattttttg	ggctggagag	agaggcccg	420
agggtagtg	aggggtgggg	tgaggagaga	gagcgagaaa	gagtgagag	agaaatttga	480
tatgaaatcc	tacatatatt	acagattgta	atgttctaaa	ctataacgat	ttgtcataaa	540
cacatatcat	ggatttgtct	ttttgtgtaa	ttttcccaat	tgtaaatagg	acttcgttat	600
ttgaaacttg	aaagtgaagt	cacatagatt	aagtacaaac	attaattaaa	gaccgtgggt	660
gaatgataaa	tattttattta	tctttaatta	gttatttttt	tgggagctct	ttattccaat	720
gtgagacttt	tgcgacatat	attcaaat	aatcgaatca	caatatgtat	tagattgata	780
aaaaaataat	ttttttacaa	tgtaggttga	gactcataac	ttactgccta	ttggtaatct	840
atgactccta	attccttaat	tatttaata	tatcatcttg	atcggttaaca	aagtaatttc	900
gaaagaccac	gagtaagaag	acaaacgaga	ataccaaaaa	attcaaaaat	ttaatgtgat	960
ttggtcaatc	gatctacgtc	cataaaggag	atgagtaatc	tactataaat	atgagagtac	1020
aaaatacaga	gagaaacaac	ctcaactaat	tcactcggaa	tacatgagaa	gttcacacaa	1080
gtgataacgt	atcaaaactg	tgaccacac	ttttccctct	aaccaaagct	cttaaaaacta	1140
tattgtgaat	gctgattaag	ttaaacgaaa	cagtcctaaa	tcttttccgt	cctatgagaa	1200
acaagattaa	tcaattcaca	atttttttta	aaagaaaaac	ctgtaagaaa	tttaggcaaa	1260
caaaacctaa	cacaagtttg	tttttgtttt	tactaccaac	aagaaattca	aatggcaaat	1320
gtataacgca	tcttagctaa	ttatatgacc	agattcagat	taatatacat	cttcacccat	1380
gcaatccatt	tctatataaa	gaaacataca	cgaacttgat	attattagag	attgagcaat	1440
ggagggtaac	aacagcagta	gcaagtcaac	caccaatcca	gcattggatc	cggatctgga	1500
cagcccggt	cagccgggtc	tggagtttgc	ccaatttgct	gccggctgct	tttggggagt	1560
cgaattggct	ttccagaggg	ttggaggagt	agtgaagacg	gagggtgggt	actctcaggg	1620
gaatgtccat	gacccgaact	acaagcttat	ttgtccgga	acaaccgaac	atgccgaggg	1680
cattcggtac	cagtttgacc	cgaatgtctg	cccgtattcc	aatctccttt	ctctattttg	1740
gagtcgccat	gacccgacca	ctctaataat	ccaggatca	aattcctttg	gtgtttcatt	1800
ttatgtgatt	aatattaaaa	attttttata	taaatgtcat	gatgatgggt	gttgctaggg	1860
taatgtagtg	ggaaagcaat	accgctcagg	aatatattac	tataatgatg	ctcaggctca	1920
actggcaagg	gagtcgtag	aagctaagca	gaaggaattt	atggataaga	aaattgtcac	1980
tgaatttctt	cctgctaaga	gatttttatag	agctgaagag	tatcaccagc	aatatctaga	2040
gaaggggtggg	ggcagaggtt	gtaagcagtc	ggctgcaaa	ggctgcaatg	acccaataag	2100
gtgctacggt	tgacagcaga	tctttgaatg	tcatagcaac	tacaaaagaa	cttgtagac	2160

```

atttgctgtc ttgcttcttt aaatttgaat aaacatgaca atgattctta taactacttg 2220
ctctcttgga tggaataaact agttgtcgta aagtattctc ctcttgctaa ttattatctc 2280
tctttatatg gtacctgcaa tttgttgctt tagttacaga ataatggacg tcaattctat 2340
atcttaattt gttttaagtc ttaaatagagg tggtttggtg ttgaaagcaa tatcaagcat 2400
agtaatacca atgatttagt agatgaactt aatcaaatca aattccaaaa tgcagtctac 2460
aaattgacaa catgaagtta agtgtatctt atgtaaattg acatctttcc tagtagatgc 2520
ctaatacttt tgtaaagact aaaataagca cagatgaggc ttgtgcattt aacttagagt 2580
tcatecttag gtgtggctgc aggagaccct gtaggggtgc ttgaagtctt gatggggtag 2640
gagggttgca ttgctatacc acacaacccc tcttcagcgt caaccttgcg ctgcattcta 2700
atgtatcctt tttctcccca ttcagctccc catgagttct tcacaatcca gtatttggtt 2760
ccatcgacgg ttgtgccata ccccaacaata gccaca 2796

```

&lt;210&gt; 9

&lt;211&gt; 363

&lt;212&gt; PRT

&lt;213&gt; Solanum lycopersicum

&lt;400&gt; 9

```

Met Glu Ser Pro Arg Val Glu Glu Ser Tyr Asp Lys Met Ser Glu Leu
  1          5          10          15
Lys Ala Phe Asp Asp Thr Lys Ala Gly Val Lys Gly Leu Val Asp Ser
          20          25          30
Gly Ile Thr Lys Val Pro Gln Ile Phe Val Leu Pro Pro Lys Asp Arg
          35          40          45
Ala Lys Lys Cys Glu Thr His Phe Val Phe Pro Val Ile Asp Leu Gln
          50          55          60
Gly Ile Asp Glu Asp Pro Ile Lys His Lys Glu Ile Val Asp Lys Val
          65          70          75          80
Arg Asp Ala Ser Glu Lys Trp Gly Phe Phe Gln Val Val Asn His Gly
          85          90          95
Ile Pro Thr Ser Val Leu Asp Arg Thr Leu Gln Gly Thr Arg Gln Phe
          100          105          110
Phe Glu Gln Asp Asn Glu Val Lys Lys Gln Tyr Tyr Thr Arg Asp Thr
          115          120          125
Ala Lys Lys Val Val Tyr Thr Ser Asn Leu Asp Leu Tyr Lys Ser Ser
          130          135          140
Val Pro Ala Ala Ser Trp Arg Asp Thr Ile Phe Cys Tyr Met Ala Pro
          145          150          155          160
Asn Pro Pro Ser Leu Gln Glu Phe Pro Thr Pro Cys Gly Glu Ser Leu
          165          170          175
Ile Asp Phe Ser Lys Asp Val Lys Lys Leu Gly Phe Thr Leu Leu Glu
          180          185          190
Leu Leu Ser Glu Gly Leu Gly Leu Asp Arg Ser Tyr Leu Lys Asp Tyr
          195          200          205
Met Asp Cys Phe His Leu Phe Cys Ser Cys Asn Tyr Tyr Pro Pro Cys
          210          215          220
Pro Gln Pro Glu Leu Thr Met Gly Thr Ile Gln His Thr Asp Ile Gly
          225          230          235          240
Phe Val Thr Ile Leu Leu Gln Asp Asp Met Gly Gly Leu Gln Val Leu
          245          250          255
His Gln Asn His Trp Val Asp Val Pro Pro Thr Pro Gly Ser Leu Val
          260          265          270
Val Asn Ile Gly Asp Phe Leu Gln Leu Leu Ser Asn Asp Lys Tyr Leu
          275          280          285
Ser Val Glu His Arg Ala Ile Ser Asn Asn Val Gly Ser Arg Met Ser
          290          295          300
Ile Thr Cys Phe Phe Gly Glu Ser Pro Tyr Gln Ser Ser Lys Leu Tyr
          305          310          315          320
Gly Pro Ile Thr Glu Leu Leu Ser Glu Asp Asn Pro Pro Lys Tyr Arg
          325          330          335
Ala Thr Thr Val Lys Asp His Thr Ser Tyr Leu His Asn Arg Gly Leu

```

340 345 350  
 Asp Gly Thr Ser Ala Leu Ser Arg Tyr Lys Ile  
 355 360  
 <210> 10  
 <211> 196  
 <212> PRT  
 <213> Solanum esculentum  
 <400> 10  
 Met Glu Gly Asn Asn Ser Ser Ser Lys Ser Thr Thr Asn Pro Ala Leu  
 1 5 10 15  
 Asp Pro Asp Leu Asp Ser Pro Asp Gln Pro Gly Leu Glu Phe Ala Gln  
 20 25 30  
 Phe Ala Ala Gly Cys Phe Trp Gly Val Glu Leu Ala Phe Gln Arg Val  
 35 40 45  
 Gly Gly Val Val Lys Thr Glu Val Gly Tyr Ser Gln Gly Asn Val His  
 50 55 60  
 Asp Pro Asn Tyr Lys Leu Ile Cys Ser Gly Thr Thr Glu His Ala Glu  
 65 70 75 80  
 Ala Ile Arg Ile Gln Phe Asp Pro Asn Val Cys Pro Tyr Ser Asn Leu  
 85 90 95  
 Leu Ser Leu Phe Trp Ser Arg His Asp Pro Thr Thr Leu Asn Arg Gln  
 100 105 110  
 Gly Asn Asp Val Gly Lys Gln Tyr Arg Ser Gly Ile Tyr Tyr Met  
 115 120 125  
 Asp Ala Gln Ala Gln Leu Ala Arg Glu Ser Leu Glu Ala Lys Gln Lys  
 130 135 140  
 Glu Phe Met Asp Lys Lys Ile Val Thr Glu Ile Leu Pro Ala Lys Arg  
 145 150 155 160  
 Phe Tyr Arg Ala Glu Glu Tyr His Gln Gln Tyr Leu Glu Lys Gly Gly  
 165 170 175  
 Gly Arg Gly Cys Lys Gln Ser Ala Ala Lys Gly Cys Asn Asp Pro Ile  
 180 185 190  
 Arg Cys Tyr Gly  
 195

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/82, 5/10</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 99/14316</b> <b>(43) International Publication Date:</b> 25 March 1999 (25.03.99)
<b>(21) International Application Number:</b> PCT/US98/19571 <b>(22) International Filing Date:</b> 18 September 1998 (18.09.98)  <b>(30) Priority Data:</b> 60/059,234 18 September 1997 (18.09.97) US  <b>(71) Applicant (for all designated States except US):</b> AGRITOPE, INC. [US/US]; 16160 S.W. Upper Boones Ferry Road, Portland, OR 97224-7744 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> BESTWICK, Richard, K. [US/US]; 6680 S.W. Canby, Portland, OR 97223 (US). KELLOGG, Jill, A. [US/US]; 6680 S.W. Canby, Portland, OR 97223 (US).  <b>(74) Agents:</b> JUDGE, Linda, R. et al.; Dehlinger & Associates, P.O. Box 60850, Palo Alto, CA 94306-0850 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>  <b>(88) Date of publication of the international search report:</b> 10 June 1999 (10.06.99)
<b>(54) Title:</b> SYNTHETIC HYBRID PLANT PROMOTER  <b>(57) Abstract</b>  The present invention is directed to a synthetic hybrid promoter composed of polynucleotide segments derived from the E8 and E4 gene promoters. The hybrid promoter is capable of providing high-level expression of heterologous genes, particularly in transformed fruit. DNA constructs containing the E8-E4 hybrid promoter operably linked to an exemplary heterologous SAMase gene are effective in conferring a delayed ripening phenotype to transformed fruit.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 98/19571

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 6 C12N15/82 C12N5/10

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 35387 A (EPITOPE INC) 28 December 1995 see the whole document ---	1-24
A	WO 94 24294 A (EPITOPE INC) 27 October 1994 see the whole document ---	1-24
A	WO 95 35388 A (EPITOPE INC) 28 December 1995 see the whole document ---	1-24
A	WO 91 01373 A (CALGENE INC) 7 February 1991 see the whole document ---	1-24
A	WO 92 17597 A (ICI PLC) 15 October 1992 see claims 1-6 ---	1-24
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

**\* Special categories of cited documents :**

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "A" document member of the same patent family

Date of the actual completion of the international search

13 April 1999

Date of mailing of the international search report

27/04/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Maddox, A

## INTERNATIONAL SEARCH REPORT

onal Application No

PCT/US 98/19571

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	XU, R., ET AL.: "Ethylene control of E4 transcription during tomato fruit ripening involves two cooperative cis elements" PLANT MOLECULAR BIOLOGY, vol. 31, 1996, pages 1117-1127, XP002099596 see the whole document ---	1-24
A	MONTGOMERY, J., ET AL.: "Identification of an ethylene-responsive region in the promoter of a fruit ripening gene" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA., vol. 90, July 1993, pages 5939-5943, XP002099597 WASHINGTON US see the whole document ---	1-24
A	DEIKMAN, J., ET AL.: "Organization of ripening and ethylene regulatory regions in a fruit-specific promoter from tomato (Lycopersicon esculentum)" PLANT PHYSIOLOGY, vol. 100, 1992, pages 2013-2017, XP002099598 see the whole document ---	1-24
A	WO 92 11371 A (ICI PLC) 9 July 1992 see the whole document ---	19,23
A	AYUB R ET AL: "EXPRESSION OF ACC OXIDASE ANTISENSE GENE IN MELON" PLANT PHYSIOLOGY, vol. 108, no. 2, June 1995, page 150 XP002034809 see the whole document -----	19,23

# INTERNATIONAL SEARCH REPORT

information on patent family members

Initial Application No

PCT/US 98/19571

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9535387	A	28-12-1995	US 5750864 A	12-05-1998
			AU 8127594 A	15-01-1996
			CA 2195095 A	28-12-1995
			EP 0766742 A	09-04-1997
			SG 50638 A	20-07-1998
			US 5859330 A	12-01-1999
WO 9424294	A	27-10-1994	US 5723746 A	03-03-1998
			AU 690547 B	30-04-1998
			AU 6530994 A	08-11-1994
			CN 1121356 A	24-04-1996
			EP 0693127 A	24-01-1996
			JP 8509122 T	01-10-1996
			SG 49118 A	18-05-1998
			US 5859330 A	12-01-1999
WO 9535388	A	28-12-1995	US 5750870 A	12-05-1998
			AU 2705995 A	15-01-1998
WO 9101373	A	07-02-1991	EP 0409625 A	23-01-1991
WO 9217597	A	15-10-1992	AU 1437892 A	02-11-1992
WO 9211371	A	09-07-1992	AU 660637 B	06-07-1995
			AU 9105691 A	22-07-1992
			AU 660638 B	06-07-1995
			AU 9106191 A	22-07-1992
			CA 2098840 A	21-06-1992
			CA 2098841 A	21-06-1992
			EP 0563157 A	06-10-1993
			EP 0563161 A	06-10-1993
			WO 9211372 A	09-07-1992
			JP 6503953 T	12-05-1994
			JP 6503954 T	12-05-1994
			US 5484906 A	16-01-1996
			US 5449764 A	12-09-1995